


SF
489
B2
P96
6th ed.

Bantams

as a Hobby



By P. Proud

With Additions by H. INMAN

ALBERT R. MANN
LIBRARY
NEW YORK STATE COLLEGES
OF
AGRICULTURE AND HOME ECONOMICS
AT
CORNELL UNIVERSITY



THE GIFT OF
PAUL POMEROY IVES 2D
IN MEMORY OF
PAUL POMEROY IVES

[illegible]

Bale
SF
489
B2
P96
6th ed.



3 1924 003 110 917

man:

BANTAMS AS A HOBBY

BY
P. PROUD.

88

SIXTH
EDITION.

REVISED
BY
H. INMAN.



FRIZZLES.	SEBRIGHTS.	
	ROSECOMBS.	
HAMBURGHES.	SILKIES,	JAPANESE.
SPANISH.	which figure in the centre, are not	POLISH.
	now classified as Bantams, but are	WYANDOTTES.
	invaluable as brooders, either pure	
	or cross-bred.	
INDIAN GAME.	MODERN GAME.	SCOTCH GREYS.
PEKINS.		MALAYS.
		OLD ENGLISH GAME.
		BRAHMAS.

"THE FEATHERED WORLD,"

9, ARUNDEL STREET, STRAND, LONDON, W.C..

HCT

SF
489
B2
P96
6th ed

Balcony

E 6771

PREFACE TO FIRST, SECOND AND THIRD EDITIONS

HAVING been frequently asked by young fanciers for information respecting the successful breeding and management of that most beautiful and profitable miniature fowl, the Bantam, I trust that the following practical information and advice will enable those who have hitherto failed in their efforts, to make their hobby both a pleasurable and profitable one. Bantams, kept under proper management, and by following the advice in mating and rearing which are fully treated of in this book, should with a little care and attention be made to not only pay the rent of a cottage and three or four acres of land, but in addition yield a nice profit and furnish a very interesting and healthful hobby as well. My readers will scarcely credit the fact that there are in England to-day several Bantam breeders who are making an income of from £75 to £250 on Bantams alone. This is not to be wondered at when we know that prices are obtained ranging from £15 to £50 for really first-class exhibition specimens, whilst the carriage to and from shows is quite 50 per cent. less than on larger breeds, and the cost of feeding is proportionately smaller. Still, it is not all "Klondyke" with Bantam farming—far from it. The young fancier will have to experience innumerable troubles and worries ere he reaches the top rung in the ladder of Bantam fame. His watchword should be, "Attention and perseverance."

In writing these notes, which are intended for the amateur fancier and not the professional, I shall endeavour to supply the necessary information in as simple a form as possible. All the varieties are treated of in their turn, according to their popularity. Full particulars are given of their respective exhibition properties, with instructions how to mate and rear, and as to their general treatment in sickness and in health. The information contained herein is not the outcome of literary knowledge, but of practical experience, and every detail is given that may be of assistance to those who are desirous of taking up a hobby which will give both pleasure and profit in return for a small outlay and a little careful attention daily.

P. PROUD.

PREFACE TO SIXTH EDITION

AN entirely new edition of this work being necessary, its enlargement and revising has again been entrusted to Mr. H. Inman, who has added notes on the varieties which have been evolved since the previous editions, amended others, and thoroughly revised the entire work.

This book was first written by Mr. Proud, and his death in October, 1911, was a real loss to poultry keepers and the fancy generally. For many years he had been chief poultry expert to *The Feathered World*, and by his wise and helpful counsel had rendered yeoman service to the cause of poultry culture.

Bantams, as being Mr. Proud's earliest favourites, always held a warm place in his heart, and it is gratifying that his book devoted to their interests has met with such appreciation as to make a further large edition necessary. The revision of Mr. Proud's standard work once more by Mr. H. Inman, his friend and successor as poultry expert to *The Feathered World*, has brought the book thoroughly up to date, and will, we trust, prove of service to all interested in the development of the Bantam fancy.

THE EDITOR.

CONTENTS

	PAGE		PAGE
Anconas - - - - -	74	Minorca - - - - -	59
Andalusians - - - - -	59	" scale of points - - - - -	59
" scale of points - - - - -	60	Nankins - - - - -	50
Baskets for Exhibition - - - - -	60	" scale of points - - - - -	50
Belgians - - - - -	68-70	Orpingtons, buff - - - - -	58
Booted - - - - -	44-46	" scale of points - - - - -	58
" scale of points - - - - -	46	" black - - - - -	73
Brahmas - - - - -	36-38	O.E. GAME - - - - -	23-26
" scale of points - - - - -	38	Pekins - - - - -	30-36
Cleaning - - - - -	81	" blacks - - - - -	33
Cochins or Pekins - - - - -	30-36	" buffs - - - - -	32
" Blacks - - - - -	33	" cuckoos - - - - -	34
" Buffs - - - - -	32	" partridge - - - - -	35
" Cuckoos - - - - -	34	" scale of points - - - - -	36
" Partridge - - - - -	35	" whites - - - - -	34
" Scale of points - - - - -	36	Perches - - - - -	59
" Whites - - - - -	34	Polands - - - - -	52
Coops - - - - -	74	" scale of points - - - - -	53
Cuckoos or Scotch Greys - - - - -	53-55	Preparing for exhibition - - - - -	83
" scale of points - - - - -	55	Rhode Island Reds - - - - -	71
Diseases - - - - -	85-87	Rocks, barred - - - - -	55-56
Dubbing - - - - -	79	" scale of points - - - - -	56
Dust-bath - - - - -	79	" buff - - - - -	58
Faverolles - - - - -	73	" scale of points - - - - -	59
Feeding chicks - - - - -	75	Rosecombs - - - - -	26-29
" adults - - - - -	76	" scale of points - - - - -	29
" before exhibition - - - - -	84	Scotch Greys (see Cuckoos). - - - - -	
" on return - - - - -	84	Sebrights - - - - -	46-48
Frizzles - - - - -	56-58	" golds - - - - -	47
" scale of points - - - - -	58	" scale of points - - - - -	48
GAME, MODERN - - - - -	9-23	" silver - - - - -	47
" birchens and whites - - - - -	21	Silkies - - - - -	48
" black-reds - - - - -	10	" scale of points - - - - -	50
" brown-reds - - - - -	18	Spanish - - - - -	60
" duckwings - - - - -	16	" scale of points - - - - -	60
" piles - - - - -	13	Sumatra Game - - - - -	72
" whites - - - - -	21	Sussex - - - - -	72
" other colours - - - - -	23	Training and Preparing for Exhibition - - - - -	83
Hamburgs - - - - -	60-63	VARIETY BANTAMS - - - - -	26-74
" pencilled - - - - -	62	" Andalusians - - - - -	60
" scale of points - - - - -	63	" Belgian - - - - -	68
" spangles - - - - -	60	" Booted - - - - -	44
Hatching - - - - -	74	" Brahmas - - - - -	36
Houdans - - - - -	73	" Cochins or Pekins - - - - -	30
Houses and Runs - - - - -	80	" Cuckoos - - - - -	53
Housing - - - - -	78	" Frizzles - - - - -	56
Indian Game - - - - -	40-43	" Hamburgs - - - - -	60
" scale of points - - - - -	43	" Indian Game - - - - -	40
Japanese - - - - -	51	" Japanese - - - - -	51
" scale of points - - - - -	52	" Jubilee Indian Game - - - - -	43
Jubilee Indian Game - - - - -	43-44	" Leghorns - - - - -	59
Langshans - - - - -	73	" Malays - - - - -	38
Leghorns, buff - - - - -	59	" Minorcas - - - - -	59
" scale of points - - - - -	59	" Nankins - - - - -	50
Malays - - - - -	38-40	" Orpingtons, Buff - - - - -	58
" scale of points - - - - -	40	" Pekins or Cochins - - - - -	30
		" Polands - - - - -	52

CONTENTS

	PAGE		PAGE
VARIETY BANTAMS— <i>cont.</i>		Washing - - - - -	82
" " Rosecombs - - - - -	26	Water - - - - -	78
" " Rocks, barred - - - - -	55	Wyandottes - - - - -	63-68
" " " other than barred - - - - -	58	" Partridge - - - - -	64
" " Rhode Island Red - - - - -	71	" Silver Pencilled - - - - -	65
" " Scotch Greys - - - - -	53	" Blacks - - - - -	65
" " Sebrights - - - - -	46	" Whites - - - - -	66
" " Silkies - - - - -	48	" Columbians - - - - -	66
" " Spanish - - - - -	60	" Other Colours - - - - -	67
" " Sumatra Game - - - - -	72	Yokohamas - - - - -	71
" " Sussex - - - - -	72		
" " Yokohamas - - - - -	71		

LIST OF ILLUSTRATIONS

	PAGE		PAGE
Ancona Bantams - - - - -	75	Nankin Bantams - - - - -	50
Belgian Bantams - - - - -	69, 70	Old English Pullet, Brown-red - - - - -	24
Birchen Modern Game, Cock - - - - -	22	" Partridge - - - - -	26
" " " Cock and Pullet - - - - -	21	" Cock, Spangle - - - - -	25
Black-red Modern Game, Cock - - - - -	10, 11, 12, 13	" Wheaten - - - - -	24
" " " Hen - - - - -	10	Pekins or Cochins, Black - - - - -	30, 32, 35
Booted Bantams, White - - - - -	45	" " " White - - - - -	33, 34
Brahma Cockerel, Dark - - - - -	36	Pile "Game" Cock - - - - -	16
" Cock, Light - - - - -	38	" Hen - - - - -	16
" Pullet, Light - - - - -	37	" Pullet - - - - -	14, 15
Brown-red Modern Game - - - - -	20	Polish Hen, White - - - - -	52
Cochins or Pekins, Black - - - - -	30, 32, 35	" " Laced - - - - -	53
" " White - - - - -	33, 34	Rhode Island Red Cock - - - - -	71
Duckwing Modern Game, Cock - - - - -	19	Rosecombs, Black - - - - -	27, 28
" " " Cockerel - - - - -	17	" White - - - - -	29
" " " Hen - - - - -	18	Scots Grey Pullet - - - - -	54, 55
" " " Pullet - - - - -	17	Sebright Hen, Silver - - - - -	46
Frizzle and Japanese Bantams - - - - -	51	" Pullet, Gold - - - - -	47
" Hen, Blue - - - - -	57	Silkie, White - - - - -	49
" Cock, White - - - - -	77	Sussex Cock, Light - - - - -	73
Hamburgh, Golden Pencilled - - - - -	62	" Pullet, Light - - - - -	73
" Silver - - - - -	61	Variety Bantams - - - - -	31
House and Run, Lean-to - - - - -	79	Washing a Bird - - - - -	82
House and Run - - - - -	80	Wyandotte Cock, Black - - - - -	66
Indian Game, Cock - - - - -	41, 42	" Hen, Black - - - - -	65
" " Hen - - - - -	41	" Hen, Partridge - - - - -	64
Japanese and Frizzle Bantams - - - - -	51	" Pullet, Partridge - - - - -	63
" Cock, Blue - - - - -	52	" Cock, White - - - - -	67
Jubilee Indian Game, Cock - - - - -	43	" Hen, White - - - - -	67
" " " Hen - - - - -	44	" Columbian - - - - -	68
Lemon Blue Game Bantam Cock - - - - -	23	Yokohama Cock - - - - -	71
Malay Cockerel, White - - - - -	39	" Cock, Duckwing - - - - -	72
Minorca, Cock - - - - -	59		

BANTAMS AS A HOBBY

CHAPTER I

GAME BANTAMS

THESE will naturally first receive our attention, because they are at the present day, and have been for years, the most popular ; and foremost of these are the Black-reds, followed by Piles, Duckwings, Brown-reds, Birchens and Whites. The reason why Black-reds have always retained their popularity is, we believe, in the first place, because they are much easier to breed true to colour, and when properly mated very few "wasters" are produced, whilst they require much less trouble in preparing for exhibition than the Piles, which have to be washed frequently, and it is well known that continual washing spoils the plumage and robs the feather of its lustre. Never wash a bird until you are really compelled to do so, for once you start you must continue. Full particulars on washing and drying will be given in due course.

In judging Game Bantams *shape and style* should be the most important, although a nice rich colour is very pleasing to the eye. The male bird should be tall and graceful, shoulders prominent and squarely set, back short and flat (a round back being a very serious fault), tapering off towards the stern ; the thighs should be long and muscular and set well apart, shanks fine and round and as long as possible. Flat shins should be guarded against ; besides counting against the bird in the show-pen, they are a sign of weakness, and care should be taken not to breed from flat-shinned stock birds, as this defect is hereditary. The toes should be long and straight, the hind toe to be carried as near as possible in a direct line with the middle toe, the point of which should just touch the ground, although in cases where birds have a large amount of style, there will be a tendency to carry the back toe too high. This is a minor fault, but where the back toe is carried sideways or inwards this is a serious defect, and amounts to disqualification in the show-pen ;

it is commonly called "duck-footed." In some cases this is caused by improper perches, but in others it is hereditary, and should be strictly noticed when making up the breeding-pen.

The head should be long and snaky, eyes large and prominent ; in Black-reds, Piles, Duckwings, and Whites the colour of the eye should be a bright red ; a light or very dark eye in these varieties counts heavily against the bird in the show-pen. In Brown-reds and Birchens the darker the eye the better, as near black as possible. The neck should be long and fine, the hackle feathers fitting closely together, wings short and nicely curved, and just meeting at the stern, carried close up, and fitting tightly to the body. Some years ago there was a tendency for Game Bantams to be long-winged, but this fault appears to have been overcome. In some cases the bird carries his wings across the back, which is a bad fault, and is known by the name of "goose-winged."

The tail, which is a most important feature in all Game Bantams, should be small and fine, the feathers carried closely together slightly above the line of the body, but not high. The feathers of the tail proper should number fourteen, and should be narrow and fine ; the sickles or long tail feathers of the male bird should be as fine as possible, reaching from two to three inches beyond the tail, and should be carried slightly curved, but not forked.

Size will next engage our attention, and it is the most important feature, not only in Game Bantams, but in every variety of Bantams, and requires very careful consideration. At the present day it is the aim and desire of most Bantam breeders and exhibitors to produce the smallest and finest-boned specimens which, although very handsome and valuable as exhibition birds, are certainly not to be relied upon as stock birds, for nine out of every dozen of these fine specimens are quite unable to pass their first egg, and die egg-bound, but if by chance they do lay, the produce from these



BLACK-RED MODERN GAME BANTAM HEN.

Bred by, and the property of, Mr. W. Rylance.

birds are little puny mites, which it is next to impossible to rear to maturity. This is one reason why young fanciers find great difficulty in rearing bantams. If you have one or two of these small, finely-bred exhibition birds that have been doing the rounds of shows, and you contemplate breeding from them, my advice is "Don't" and keep them from laying as long as possible.

Pullets for the breeding-pen should not be less than fifteen ounces, and from two to three ounces added to this would give better results; and cockerels sixteen to twenty ounces are quite small enough for all practical purposes.


Of course proper feeding has much to do with size, and this shall receive attention in due course.

We now come to the next most important feature, viz. *colour*, which in the show-pen should count fifteen points.

BLACK-REDS

In Black-red cocks, the face, head, lobes and wattles should be a bright, healthy red, the neck hackle bright orange, free from striping; back and wing-bow rich crimson, whilst the

saddle hackle should match the neck hackle as near as possible, wing-butts black, wing-bars a glossy or steel blue, whilst the bays or wing-ends should be a light chestnut; legs, feet and beak, rich olive or willow, but not blue, which we often find in some strains where Duckwing blood has been introduced, but in the breeding-pen blue legs should be guarded against. The breast and thighs should be black, perfectly free from ticking or lacing, although in adult birds it is rather difficult to get a cock perfectly black in breast. The tail and sickle feathers should be black throughout, including the shaft which runs through the centre of the feather, and which in some strains is more or less red, and generally denotes a pullet-breeding strain; this is, of course, objectionable in the show-pen, although not to any great extent, still a black shaft is at all times preferred.



BLACK-RED BANTAM COCK, "BRIGHT 'UN."

A favourite stock bird of Mr. H. A. Morris

We will now go on to describe the exhibition Black-red pullet to match the cockerel, and which should be identically the same in colour, eyes, face, head, comb, and lobes; white in lobe being a great drawback, although it is a very difficult matter to breed pullets absolutely free from white in lobe. This blemish is sometimes removed by the professional exhibitor with a pair of scissors, an operation demanding considerable skill, and, quite apart from any justification of the act, which is certainly a very questionable one, it may be added, it can only be performed by a very steady hand, without which the remedy being worse than the defect. Honest exhibitors will do well to leave such operations alone.

The neck hackle of the pullet should be a pale golden colour with very narrow black stripes on each side of the shaft, with a light golden edge round the black. The body and wings should be one uniform shade of brownish drab, a nice soft, even colour throughout, very finely pencilled with black, this point being of great importance, as the pencilling has a tendency to be too large and coarse, a defect which is very objectionable in the show-pen; although in cockerel breeding pullets showing coarse pencilling are sometimes very useful, as such pullets generally throw cockerels with good sound bays.

The throat should be a pale salmon, breast rich deep salmon, shading lighter towards the thighs and under the body. The tail should be black, with the exception of the two outer or top feathers, which should correspond as nearly as possible with the colour of the body.

I have now described the standard colours of the Black-reds, and will now proceed to give the young fancier advice as to the best way of mating up his breeding-pens to produce as nearly as possible the necessary requirements above mentioned.

Now, in the first place, see that your breeding stock is perfectly healthy and free from disease. Never breed from unhealthy birds, as it is nothing but labour in vain. If your breeding stock is unhealthy your chicks will be 50 per cent. more so, and here let me say that the fewer birds you have in your breeding-pen the stronger will be the produce. Be content to breed a few good useful birds rather than a greater number of inferior ones, and remember that one really good bird is worth forty bad ones, either in the show-pen or for stock. Much has been said about the proper number of hens or pullets to be used in the breeding-pen to

produce the greater number of pullets and vice-versa, viz. that if eight hens or pullets are mated to a two-year-old cock, 75 per cent. of the produce will be pullets, and that if only two or three hens or pullets are put with a cock the same percentage in cockerels will be the result. If you pin your faith to the above you will be greatly disappointed, as the experiment has been frequently tried, and in nearly every instance the result has been in many cases the opposite to what has been looked for. My contention is that if a hen is to produce a greater number of pullets than cockerels she will do so, whether there be three or eight hens



WINNING BLACK-RED MODERN GAME BANTAM
COCK.

The property of Mr. R. Fletcher Hearnshaw.

in the breeding-pen, and the same applies in cockerel breeding. If the young fancier has sufficient accommodation it is best to mate up two pens, one for cockerel breeding and the other for pullet breeding.

Now, in mating up the *cockerel pen*, I should take the tallest and brightest-coloured cockerel I could find, preference being given to one that is short in back with good prominent shoulder points, and care being taken that his breast bone is straight, and that his back toe is well carried in a direct line with the middle toe. Having selected the cockerel, we should now

pick out the hens or pullets, latter preferred, to mate with him, with a view to producing good, bright-coloured exhibition cockerels. To do this we shall require three reachy tight-feathered pullets, very pale in hackle to crown of head, discarding those with dark caps and coppery hackles, which are useless for cock-breeding. The hackle should be as silvery as possible, right up to the comb, the body-colour and wings be inclined to red or foxy, with a rich golden edge round each feather. Select the pullets that are shortest in body and tail, and especially those which show a lot of style and reach, and the more prominent the shoulders are the better.

Some breeders prefer to breed from a wheaten pullet to obtain the bright top colour; and although a very satisfactory cross, and one which will give good results in cockerel-breeding, yet should this strain by any chance get mixed up with your pullet strain, you will know it to your sorrow, and it will take years to repair the mischief done. Therefore if you decide to use wheaten blood in your cockerel pen, keep the produce rigorously by themselves

and have them all carefully rung; this is the only sure road to success in breeding.

For my own part I prefer to use nothing but Black-red pullets in cockerel-breeding, such as those stated above, for the simple reason that should one bird get astray the harm done is comparatively small, and the produce will be found in every way quite as satisfactory.

The following season I should pick out three or four of the most likely-looking pullets bred from this pen, especially those with plenty of reach, the taller the better, and with those I should mate the sire of them, and put the most suitable cockerel with the hens in place of the old bird. By doing so you are building up a strain of your own, which, with care and attention, cannot fail sooner or later to reach to success, for it is a well-known fact that to secure colour and the other necessary points in bantams, you must inbreed; still, at the same time it must be borne in mind that there is a happy medium in inbreeding beyond which it is dangerous to proceed, and in this the young fancier should be guided by the health and stamina of his stock, for if inbreeding be carried too far, the produce will become weakly and difficult to rear; in that case fresh blood should be introduced, and this is best done by obtaining a male bird with a dash of the same strain in him if possible, but not closely related.

Having mated up the cockerel-breeding pen of Black-reds, I will now give attention to the *pullet pen*. In the first place, the hens or pullets must be sound in colour, free from rust or coarse pencilling; this is of the greatest importance in pullet-breeding. It is only a waste of time trying to produce exhibition pullets from hens which are faulty in colour; it is utterly impossible to do so. Besides being sound in colour, the hens must be of good shape, not too small, and the shorter in back the better. The male bird should be from a pullet-breeding strain, and in colour should be darker than the cockerel-breeder. I always prefer a bird of a brickish top colour, and one uniform shade from his hackle to tail. His wing bar and shoulder points should be perfectly free from lacing and as black as possible. This refers to cocks as well as cockerels, although the adult birds will in all probability be laced on breast, and I have often found that such birds from a pullet-breeding strain are invariably weak in their bays or wing ends—that is, the bay of the wing end does not continue to the end of the feather; this is always a good sign when selecting a cock for pullet-breeding.



BLACK-RED MODERN GAME BANTAM COCK.

The property of Mr. Harold A. Morris.

In many cases the shaft of the sickle feather in such birds will be red instead of black. If you prefer to breed from an old bird, do not mate him to more than three pullets, especially early in the year, say February, or a large percentage of unfertile eggs will be the result. When breeding from a cockerel, hens should be used, and if he is a healthy vigorous bird, you may safely put five or six with him with good results.

In mating up the Black-reds I have taken it for granted that the young fancier has plenty of accommodation, but very many have only a very limited space at their disposal, and in such cases one pen will be sufficient. Where this is so I should take a cock that answered as far as possible to my ideal of a pullet-breeder, except I should prefer him to be a light shade in top colour, the paler the better, and with him I should mate two cockerel-breeding pullets and two sound coloured pullets for pullet-breeding. From this pen I should expect to get both exhibition pullets and cockerels, though not the same percentage of winners as from separate pens; still, it can be done, although it has been repeatedly stated that to produce exhibition specimens of any variety of poultry two pens should be used. This statement is all right to those who know no better, but from personal experience I know that it is quite possible to breed winners of both sexes from one pen of stock birds when properly mated to give this result.

Still, where the young fancier has plenty of room it is advisable to use two pens, and where it is necessary to build up a strain of your own, separate pens in such cases *must* be used. Let me state here that when breeding from only one pen never by any means use wheatears; these may only be used exclusively in cockerel pens.

PILES

The next most popular breed are *Piles*, the cockerels being the most handsome of all varieties. Although quite a distinct breed in themselves, it is quite possible to produce the very best by crossing, which I will show in due course.

In the first place, the Pile cockerel should be identically the same colour as the exhibition Black-red cockerel, but with this exception, that where the Black-red is black the Pile cockerel must be a sound white, viz., the breast, thighs, wing butts, wing bars, and tail; legs and beak

a rich orange yellow; willow legs and white legs being very objectionable. Although in the past willow-legged Piles could frequently be seen at the top of the prize list at our shows, at the present time a Pile with willow legs, however good he may be in other points, does not stand the slightest chance of figuring successfully in the show-pen. The face should be a bright healthy red, and when undubbed the earlobes should also be red and free from white. Although it is somewhat difficult to obtain, the rich ruby eye in Piles as in Black-reds, still they should be as red as possible. In the show pen, from a colour point of view, judges differ. Some prefer the top colour to be a very dark colour both in wing bow, back and hackles, the same as one would expect to find in a pullet-breeding Black-red cock. For my own part I much prefer the bright top colour and rich orange hackles, but, above all, a pure white breast and dark wing ends.

The breast of the Pile cockerel is a very important feature, as it is much easier to obtain



BLACK-RED GAME BANTAM.

Exhibited by Mr. W. D. Trickett.

the desired top colour than a perfectly white breast.

The Pile pullet should match the cockerel in point of colour of face, eye, legs, and beak; breast should be a rich salmon, shading off to the thighs; the body, wings, and tail should be a pale creamy-white, as free from red or foxiness as possible. Although a rich dark breast and a clear wing are most difficult to obtain, yet in the present-day competition a pullet or hen inclined to foxiness or rose-winged cannot hope to be successful in the show-pen, although, however careful you may be in the selection of your breeding-pen, to produce one pullet perfectly clear on breast and back, and a sound rich salmon breast, you will breed fifteen or twenty that are rose-winged more or less, or, what is still worse, a dark creamy colour throughout. When the foxiness on the wing is only confined to a few feathers on each side, those are often removed by the unscrupulous

or habitual faker, and such birds have been known to win repeatedly. Still, by a practical judge this irregularity can quickly be detected, and in such cases the bird should not be passed, but disqualified, for when these feathers grow again the bird is useless as a show specimen, unless the same operation is resorted to, which I am sorry to say is very often the case. I will now go on to show how to breed Piles, and will mate up two pens as in Black-reds.

In mating up the breeding-pens of Piles, I will in the first place endeavour to give my readers the best advice in mating up pure bred Piles—that is, birds that are bred from Piles on both sides, as many of the very best exhibition Pile cockerels are bred from Black-red cocks, and it is very important that the young fancier should know when mating up his breeding-pens how his stock birds have been produced.

Taken for granted that the Pile cock selected is the most suitable for cockerel-breeding in shape and reach, as described in the cockerel-breeding Black-red pen, the next most particular point is to see that he is perfectly sound in his white, free from lacing or smokiness on breast, and, above all, possessing a sound, deep bay on wind-end, for without the latter he is no use as a cockerel-breeder. I have never known a male bird that was weak in colour on wing-ends produce a single exhibition specimen, and this is quite as essential in breeding for pullets, therefore bear in mind that a Pile cock or cockerel weak in his bays is practically of no good whatever either as a stock bird or for exhibition, however good he may be in all other points. Next select three or four tall stylish pullets, deep in breast colour, short in back and possessing plenty of reach, and showing the shoulder-points prominently, preference being given to those which are nicely rosed on the wing, but *not* creamy on wing-ends. From this pen I should expect to get sound-coloured cockerels, although the pullets would be too warm in colour for exhibition, yet as stock birds they are invaluable, and especially so if mated back to the sire. In pullet-breeding, the cock must be sound in wing-ends, clear white in his wing bars and shoulder-points; in top colour I prefer a bird of one uniform colour of the darker or brickish colour, and if laced on breast all the better. The hens or pullets to mate to this cock should be as clear as possible on wing and back, free from rosiness or creaminess, and preference being given to



From a Painting by Miss Williams.

PILE GAME BANTAM PULLET.

Bred and Exhibited by Mr. Fred. Smalley.

those possessing good salmon breasts, although it is quite possible to breed the very best pullets from hens which are pale in breast colour, providing the body colour is perfectly clear. Cockerels bred from this pen will be found too pale in colour for exhibition birds; but those which are extra good in bays and whites should be retained as stock birds for another season. Having mated up pens of pure-bred Piles, I will now go on to show how to breed exhibition specimens by crossing with Black-reds.

It is a well-known fact that in breeding from pure-bred Piles for four or five years in succession, the top colour in the cockerels is lost altogether; therefore it is necessary to introduce Black-reds to build up the colour. The best way to do this is by selecting a Black-red cockerel, extra rich in top colour, rich mahogany saddle, the more colour the better, the breast and wing bars a sound black, free from ticking or lacing; this is a very important point if we would have Piles with a sound white breast. Another special point to note is that the wing-ends are a dark chestnut, sound from wing bar right to the very end of the wing. Do not breed from a Black-red which fails in wing-ends, as it will be only labour in vain; but if care is used in only breeding from a bird as described, and mating him to two or three white pullets, you will be rewarded by breeding the very best coloured Pile cockerel it is possible to produce. When I say white pullets I mean what are sometimes called lemon Piles—viz. Pile pullets clear on wing and almost white on breast—the clearer on breast the better. Although the cockerels from this pen will be yellow-legged, quite 75 per cent. of the pullets will be willow-legged, and as such are useless in the show pen, though generally these are clearer in body colour, more stylish, and sounder in breast than the yellow-legged pullets, and if used judiciously in mating them to a reliable pullet-breeding cock that is pure Pile bred, good-coloured pullets can be obtained.

If the Pile pullets have a tendency to come pale in breast, it would be advisable the next season to put these to a yellow-legged Black-red cockerel that has been bred from Piles, preference being given to one of a light brickish top colour of the same shade from neck to saddle hackle, care being taken that his wing bars and wing butts are free from lacing, although a little lacing on breast is no detriment whatever. To those whose space is



PILE GAME BANTAM PULLET.

Bred by, and the property of, Mr. W. Rylance.

limited to one breeding pen I should use a tall, short-bodied cock or cockerel, approaching as near as possible to the exhibition Pile cockerel in colour, and to him I should mate up two pullets or hens clear on wing and with good deep salmon breasts, and a couple of stylish and reachy rose-winged pullets, and from this pen I should look for good all-round cockerels, also pullets that would be able to hold their own in second-rate competition, although I should hardly expect the pullets to be as clear on wing as those specially bred from a pen mated up for pullet-breeding.

Piles are best kept on a good grass run, and the nature of the soil has much to do with the leg colour. A heavy clay or a light sandy soil is best for preserving the rich orange colour, while dry limestone soil will quickly transpose the rich yellow to a white. I have proved this over and again. Feeding is a very important point in Piles. Maize, although beneficial to leg colour, will have a tendency to make the body colour creamy, therefore use maize very sparingly in feeding Piles. Those Piles which are bred from Black-reds will invariably be found to have the richest leg colour, and this is



Photo]

[*"Feathered World."*]

PILE MODERN GAME BANTAM COCK.

Exhibited by Mr. Walter Firth.

a capital guide to the young fancier in introducing a strain of Black-red. As soon as he finds that the leg colour is deteriorating, it is a proof that a cross is required to strengthen the colour, although there is no fixed rule as to how long a time should elapse before introducing the Black-red, some strains requiring a change more frequently than others. Still, as a guide once in three or four years will be found to be sufficient.

Whilst on the subject of Piles, it will be as well for me to inform the would-be fancier of this very charming variety that in all the varieties of Game Bantams no other breed is so difficult to breed true to colour as Piles, and none that requires so much attention in preparing for exhibition; for to be successful in the show pen with Piles they require washing very frequently, and be it known that constant washing takes all the nature out of the feather. Fanciers who have not a good grass run with a good loamy or clay soil away from the smoke of factory chimneys, should not entertain the idea of joining the ranks of Pile fanciers. Again, breeding Pile Bantams is not all lavender—far from it. Should you be successful in rearing, say, 150 chicks, and out of this number you can manage to select twenty good enough to win,

you may count yourself one of those lucky individuals that Dame Fortune has smiled kindly upon.

There is always a good demand for really high-class exhibition Piles. Really good birds are always in demand from £5 to £15; therefore one of the latter will cover the cost of keep of a good number of wasters, which will always put in an appearance, no matter how much care you exercise in mating up your breeding-pens.

DUCKWINGS

Duckwings come next on the list in point of favour, and are a very pretty and taking variety, and, like the Piles, are rather difficult to breed true to colour. Unlike the Duckwing Leghorns, which are divided into Golds and Silvers, and exhibited as such, the Duckwing Bantam is restricted to Golds, as the Silvers



Photo]

[*"Feathered World."*]

MODERN PILE GAME BANTAM HEN.

The property of Mr. R. S. Marsden.



DUCKWING MODERN GAME BANTAM PULLET.

The property of Mr. A. Tyrie.

have not yet found favour in the exhibition world.

The Duckwing cock should, like the Black-red, be red in face and eye, breast sound black, also wing butts, wing bar and thighs down to the hock black, the neck hackle and saddle hackle a silvery white, as clear from black striping as possible. Although it is somewhat difficult to breed out this defect, it is a most important point, and should receive careful attention when mating up the breeding-pens. The saddle and wing-bow should be a rich yellow or orange colour, shading to silvery white in hackle, the wing-ends from the black wing bar to the end of the wing clear white, free from rusty colour at the outside edge of the top feathers. Quite 50 per cent. of Duckwing cockerels fail more or less in this respect. The tail of the Duckwing cock should be identically the same as described in the Black-red.

The Duckwing pullet should, like the Black-red, be red in face and eye, legs willow, her body colour and wings a light slate colour, or grey evenly and finely pencilled with black, and free from shaftiness, some strains showing more or less the white shaft on each feather, which should not be; the colour should look one even soft grey throughout, the more uniform the better.

The breast should correspond with the Black-red pullet, viz., it should be a dull salmon, although it is one of the most difficult points in Duckwing breeding to get a sound breast colour, together with a perfect body colour; the weakness being to run pale in breast, more or less. This is not a very serious defect unless the breast be very pale or almost devoid of salmon shading, then it counts heavily in the show-pen.

As Duckwings are derived from Black-reds, it is absolutely necessary that you should know



DUCKWING MODERN GAME BANTAM COCKEREL.

Exhibited by Mr. A. Tyrie.



DUCKWING MODERN GAME BANTAM HEN.

The property of Mr. R. Fletcher Hearnshaw.

how the birds you intend using in the breeding-pens have been produced.

If the Duckwing pullets are pure I should take two or three and put to a Black-red cockerel, which I have before described in the Black-red cockerel-breeding pen, and from these I should expect good coloured Duckwing cockerels but no pullets; the pullets would be Black-reds, and, being bred from the Duckwing hens, they should be marked or rung to prevent them being used in the Black-red pens another season. The best use for such pullets, if they are sound and level in colour, and clear in hackle to crown of head—i.e., free from dark in cap—is to put them to a silver Duckwing cockerel, and by doing so good-coloured Duckwing pullets should be the result. Another way to breed Duckwing pullets is by mating a sound-coloured Duckwing cock—that is, pure Duckwing bred for two or three years back, and with him mate three or four of the soundest-coloured hens or pullets, care being taken that they are not pale in breast. Any pullets from this pen that are not up to exhibition standard in point of colour should be put back to a Black-red cockerel the next season, for the purpose of breeding Duckwing cockerels.

In breeding Duckwings, where other varieties of Game Bantams are kept, it is advisable to

keep them rigorously by themselves, and their produce carefully rung, or incalculable harm can soon be done, especially amongst Black-reds.

BROWN-REDS

Brown-reds will next engage our attention. Although 50 per cent. of the shows provide classes for Black- or Brown-reds, it is very rarely we find one single specimen of the latter. What is the reason? Well, in the first place, Black-reds have always taken the lead, and I am firmly of the opinion that they always will, although the Brown-reds appear to be making steady progress, so much so that, in pullets especially, there are to-day some that could hold their own against the best Black-reds, both in type and quality; still, the judges have always hitherto given preference to Black-reds. In cockerels the Brown-reds are still a long way behind the Black-reds. The colour has been obtained almost to perfection, but there is yet too much feather, and, with one or two exceptions, the type and stateliness of the miniature Game Fowl, which we find in Black-reds, Piles, and Duckwings, are still wanting more or less in Brown-reds. Another reason why so few Brown-reds are found competing against Black-reds is that, as yet, the former are practically in the hands of three or four exhibitors, whom we find clearing the boards at our more important fixtures; and until they become more widely distributed this state of things will continue. But the main reason, I believe, is that the judges have a decided preference for the Black-reds; that is why we very seldom find Brown-reds competing. In my own mind, I think that Black-reds should compete alone, and let the Brown-reds come under the head of A. O. V. Game Bantams where classes cannot be provided for them. I believe this would in some measure be the means of making the Brown-reds more popular.

Brown-reds, like the Black-reds, breed true to colour, and are much less trouble in preparing and keeping in show form than Piles or Duckwings, and can be kept almost anywhere, where it would be next to impossible to keep the former. The Brown-red cock should have a very dark mulberry face, almost approaching black, the darker the better; the eye should be as dark as possible, a red or light eye counting heavily against the bird in the show-pen; the legs and feet should be almost black. The

neck and saddle hackle should be a pale lemon, the back and saddle rich lemon, breast black ground colour, with pale lemon lacing round each feather and showing the shaft of the feather, the lacing to start at the throat and continue nicely down the breast, but not much beyond the shoulder points. This is in direct contrast to what prevailed some years ago, when the lacing had to continue down to the top of the thighs. The curtailment of the lacing is a great improvement, for without that heavy lacing that was customary a smarter appearance is given to the bird. Wing-bar and tail black, the latter as fine and as short as possible, although it is very rarely that the Brown-red is so fine in feather as the Black-red, especially when the bird possesses that pale lemon top colour which is so much desired. A few years ago the lemon top colour had hardly been obtained; in its place we found then dark orange or nearly red, and I believe it is by breeding for colour alone that Brown-red fanciers have lost the hardness of feather; for it is a well-known fact that the lighter the colour the softer the feather, and vice versa; yet, to be successful in the show-pen, a Brown-red cockerel must possess that beautiful lemon top colour which to-day is more than half the battle, irrespective of hardness of feather. Still, I contend that in judging Game Bantams shape and type should be the first requirements, then colour and hardness of feather.

The Brown-red pullet to match the cockerel should be identically the same in face, eye and legs. The neck hackle should be a pale lemon with a narrow black stripe, and this colour should reach to the crown of the head; the weakness in Brown-red pullets is to be coppery or dark-capped, viz., the top of the hackle towards the head runs a dirty coppery colour, which is a great objection; although it is one of the mysteries of Game breeding to produce clear lemon-hackled pullets free from lacing on the back and shoulders, still, it can be done with careful mating and perseverance. The body and wing should be a glossy black, free from lacing or shaftiness; the ground colour of the breast should be black, each feather from the throat to the breast distinctly laced with pale lemon, the lacing to be uniform, and continued well down to the thighs, the breast lacing being one of the most important points, as quite 30 to 40 per cent. fail more or less in this respect.

In mating up Brown-reds for producing exhibition cockerels it is necessary to have a

separate pen, and to do this I should select the most perfect coloured male bird I could find, one possessing that beautiful pale lemon top colour and plenty of it. I should like him to have all the breast lacing possible, but at the same time I should like the lacing to be clear and distinct, and yet possessing black shoulder butts free from lacing. Another important point to look for is a sound dark eye, the blacker the better; never breed from a light-eyed bird. The pullets to mate with this cock should be as tall as possible, tight in feather, and as pale in neck hackle as possible, care being taken to see that the pale lemon is continued to the top of the head; this is one of the great secrets to obtain bright-coloured cockerels. If you want bright-coloured cockerels, never use dark-capped pullets or hens; see that they are clear lemon right up to the comb; never mind if they are laced on the back, all the better, as you will get more colour in your cockerels, and although the pullets from this pen will be useless as show specimens, they are invaluable as cock breeders, and should some of them come with bright, pale hackles, laced on back, and continued on to the wing or saddle, guard these in the same way as you would guard a Klondike claim, for there is gold in them as cockerel breeders, although in other respects they are worthless.



DUCKWING MODERN GAME BANTAM COCK.

The property of Mr. R. Fletcher Hearnshaw.

The next season it would be advisable to put these pullets back to the sire, and put the best cockerel of that season's breeding back to the hens. In breeding for pullets it is not necessary to have the cock bird so pale in colour as the cockerel breeder. One of the orange shade, or even darker, would be preferable; care must be taken to see that he is sound in eye, very clear in lacing on breast, and that the lacing is carried well down the breast, and not patchy or irregular. To him mate four or five pullets or hens that are perfectly sound in body colour, quite free from lacing on back, yet well laced on breast. This is absolutely necessary in pullet breeding. The hackle should be as clear almost and as bright as the pullets I have already described; in fact, pullets for pullet breeding should approach as near as possible to the exhibition standard; from this pen I should look for the very best coloured pullets, but the cockerels would be found deficient in colour, and of a darker shade; these, however, although only suitable for second-rate competition, should always be retained as pullet breeders, and by doing so you are building up your own strain, which is very desirable in Game Bantam breeding.

Having described the mating-up of the breeding-pens of pure Brown-reds, I will now go on to show how the young fancier may produce exhibition specimens by introducing a cross of another variety.

Years ago, when the now fashionable pale lemon top colour was first produced, many fanciers were at a loss to know how it had been obtained, although had they been able to have got to the right side of some of our celebrated large game breeders of this variety, they would have learned the secret much earlier, for the bright lemon colour had been secured in the large game years before we ever saw it in the Bantam.


How was it obtained? Simply by crossing with Birchen blood. The question that will naturally rise in the mind of my readers now will be, Yes, but how are Birchens produced? But for answer you must "bide a wee." This information will come in due course, but sufficient for the day is the evil thereof. Although I am not in favour of the introduction of Birchen blood in Brown-reds, for the simple reason that by doing so, to a certain extent, you are losing the dark gipsy face so characteristic in the pure bred Brown-red, still there are times when the young fancier

is driven into a corner as it were, and will puzzle his brains how it would be possible to get the desired pale lemon top colour and lacing already described, seeing that he does not possess the colour in the cockerel breeder.

The only remedy for this defect is to select a Birchen cock or cockerel will plenty of top colour, as silvery white as possible, and well-laced breast, and put him to three or four pullets or hens; if you use a cock then put pullets, and vice versa. In this case it is not absolutely necessary to have the hens so pale in hackle as described in the cockerel-breeding pen above.

From these you should obtain cockerels quite as bright in lemon as those from the pure cockerel pen, with the only exception that there will be a tendency for the produce to be lighter in eye and face.

From these cockerels it would be advisable to select two or three of the tallest, palest in top colour, preference being given to those having plenty of distinct lacing on breast. By using one of these with the same hens the following season, and another with the best coloured and tallest pullets, you will still retain the desired colour and at the same time be improving the face colour and eye. It would not be found advisable to use the Birchen cock a second season, unless it was with the idea of breeding more Birchens, and this could be brought about by putting him to the Birchen pullets which are certain to be found among the produce of the Birchen cock and Brown-red hens, and of course the following season the percentage of Birchen would be much greater.



BROWN-RED MODERN GAME.


BIRCHENS AND WHITES

No variety of Bantam, except Old English has grown more in public favour during the past few years than Birchens, and this is not to be wondered at, for I know of no prettier or more taking bird than the Birchen pullet.

Birchens have been bred and exhibited for many years, but it was long before they were able to make headway, and now that

retained, although it is a difficult point to obtain, the same depth of dark mulberry face colour in Birchens that is found in Brown-reds, simply because the variety has been obtained by a cross with a red-faced variety. Still, with perseverance and careful mating to breed out the defect it can be done, and has already been accomplished by one or two who have made this variety a speciality.

The Birchen pullet, in body colour, face,



MODERN BIRCHEN GAME BANTAM
PULLET.

MODERN BIRCHEN GAME
BANTAM COCK.

Both bred and exhibited by Mr. W. F. H. Butler.

they have found favour, they have every appearance of being as fully established a variety as the Duckwings and Brown-reds from which they have descended. Birchens were obtained in the first instance by crossing a Silver Duckwing cock with Brown-red hens, but at the present time they are bred true to colour from pure Birchens. The Birchen cock should resemble the Brown-red in every respect except top colour and breast lacing—viz., where the Brown-red is lemon, the Birchen should be silvery-white, the whiter the better, providing that the dark gipsy face is still

head and eye, and legs, should be identically the same as the Brown-red pullet. The neck hackle should be a silvery-white, with a narrow black stripe on each side of the white, and should be free from any other tint, such as lemon or straw, both being considered defects in Birchens, but which will be found more or less in those strains which have not been bred pure. Birchen pullets also have a great tendency to be dark-capped, i.e., the silvery hackle, which should be one distinct colour from base to the crown of the head, becomes much darker or nearly black towards the comb; this should not be,

as I pointed out in my remarks on Brown-reds, and it counts heavily against the bird in the show-pen. Another weak point in some strains of pullets is the scarcity of lacing on the breast. I have seen some of the most typical-shaped, clear hackled pullets which have been thrown back for the want of breast lacing, some years ago, but at this day the scarcity of breast lacing is not so fatal to a bird's chance in the show-pen. The breast should be well and distinctly laced from the throat to about a line about parallel with the shoulder points, and not to the top of the thigh, as was customary in years gone by. At the same time, the back and wing must be free from lacing, which is not easy to secure, as only those who have bred them know. Of course, as I have just said, the want of breast-lacing will throw a bird back considerably in the eyes of a competent judge, yet it would not be so fatal as laced back and shafty wings, providing it was as good in other points; and it would be as well for the young fancier to bear this in mind when making his selection for the show-pen.

Another stumbling block in Birchens is the colour of the eye, which should be dark brown, as near black as possible, although it is next to impossible to breed Birchens as dark in the eye as Brown-reds, but red eyes are certainly not to be countenanced, no more than a dark eye is in Black-reds, yet Birchens with red eyes are often found winning, but this is wrong, and should handicap the bird heavily in the show-pen, whilst light eyes should put the bird out of running altogether, the same as in Black-reds or Brown-reds.

In breeding Birchens, they must be mated up precisely the same as described in Brown-reds. Should the young fancier be unable to procure Birchens, which he should have little difficulty in doing by inserting an advertisement in *The Feathered World*, it would be advisable for him to get a Brown-red hen or pullet, one heavily laced on breast preferred, and mate her to a Silver Duckwing cockerel; the latter can be obtained from any breeder who makes a specialty of breeding exhibition Duckwing pullets, and such a bird can be obtained for 15s. or £1, simply because Silver Duckwing cockerels are useless in the show-pen, but essential in the breeding-pen. Still, where Birchens can be secured I strongly advise their use, in preference to crossing as above, as it is a great saving of both time and trouble.

Another variety that has come forward

more prominently during the last year or two are Whites, and, strange to say, these originally descended from Brown-reds, and at the present day some strains of Brown-reds will still throw Whites. Although a very pretty variety, I do not think they will ever "take on" with the Fancy.

Whites can also be produced by breeding from the light lemon-coloured Piles, but it will take at least three years to breed out the lemon tint in the hackle. Still, as pale lemon Piles are only "wasters" in the eye of the Pile breeder, the foundation could be laid for an outlay of 30s. for a breeding-pen. By breeding from Lemon Piles as near white as possible, the rich orange-yellow leg so necessary in Whites will be all the more readily obtained, as Whites bred from the offshoots of Brown-red are more or less willowy in leg colour, which is considered a defect in the show-pen. Plenty of these can be got, especially about August or September, when Pile breeders will be glad to dispose of such at from 7s. 6d. to 10s. each. Whites should be pure white in body colour, rich yellow legs and feet, also beak; face, comb, and eyes red. Willow legs a disqualification.



MODERN BIRCHEN GAME BANTAM COCK.
Bred and exhibited by Mr. H. L. Eaves.

Whites can only be kept satisfactorily where the atmosphere is free from smoke, and where they can have free range on a good grass run. Kept under such circumstances they look very pretty indeed.

OTHER COLOURS

Other varieties that have been seen in recent years are Blues, Blue-reds and Lemon-blues, but for some reason or other they seem to have made very little headway in the Fancy, at least as far as the number of exhibitors go. This is rather strange, as all are very pretty and interesting, and, except in colour, are identical to the other varieties of Game Bantams. A Blue-red cockerel is similar in colour of saddle, neck and saddle hackles to a Black-red cock, with solid blue breast and tail, and under parts of a rich deep blue. The pullets have blue wings and tail, and blue ground colour, with the Black-red hen colour (brownish drab) intermingling in body, and salmon breasts. A Lemon-blue cockerel has the lemon neck hackle, saddle and saddle hackle as seen in a Brown-red, paler, but sound blue under parts, tail and breast, and laced on breast with lemon like a Brown-red. The pullets of this variety are of a pretty even deep blue, lemon head and hackle, with lemon lacing on breasts as in the Brown-reds, and the rest of the colour blue. The two colours, Blue-reds and Lemon-blues, are quite distinct. In shape, style, limb and shortness of feather they should favour the other varieties of Game Bantams. They are very rare and most delicate, but still decidedly pretty, and are quite as good as any other colour of Game Bantam.

SCALE OF POINTS FOR GAME BANTAMS

Good head and neck	..	..	8
Good eyes	..	..	10
Good body and wings	..	..	10
Good legs and feet	..	..	10
Good tail	..	..	8
Good shape and style	..	..	12
Good colour	..	..	15
Good condition	..	..	8
Hardness of feather	..	..	9
Correct size, etc.	..	..	10

100

The same scale to apply to defects, and deducted accordingly.

DISQUALIFICATIONS.—Crooked breastbone, duck-footed, wry-tailed, roach-backed.

CHAPTER II

OLD ENGLISH GAME BANTAMS

No other breed of Bantam awakened such interest, or caused such a boom in the Fancy during the year 1898, as Old English. For years before this charming variety had been hanging fire, as it were, but, like all other new breeds, once they were launched in the exhibition world, and classes provided for them, their popularity was quickly assured by such a novelty lover as the English exhibitor. And why, I ask, should not the Old English



Photo]

[“ Feathered World.”

LEMON BLUE GAME BANTAM COCK.

The property of Mr. Wm. H. Green.

Game Bantam find equally as much favour as the large breed of Old English Game, from which they have descended? To-day in north-country shows and at all the big events throughout England, no classes fill better or pay better than Old English Game, and Old English Game Bantams are nearly, if not quite, as popular.


As an example of the rapid strides the Old English Bantam has made, I need only refer to Whitehaven Show of November 11th, 1898, where a Black-red cockerel which carried off first prize was claimed at catalogue price, £10, although, had this bird been seen running about in any of the back-yards twelve months previously, it might have been bought for eighteenpence. Then at Birmingham Show, three weeks later, the third prize bird was bid up to £11 11s.; and which a couple of years ago would not have fetched as many shillings.

Why is it, then? you ask. Simply a whim of the Fancy, which must have something novel, no matter what the price may be. Still,



OLD ENGLISH BROWN-RED BANTAM PULLET.

Owned by Messrs. Watson & Robinson.



O. E. WHEATEN GAME BANTAM.

The property of Mr. J. S. Appleton.

I honestly believe that Old English Bantams have come to stay. They are exceptionally hardy and healthy; they will live and thrive where it would be impossible to rear the modern Game Bantam.

The words, or shall I say opinion, of my old friend have proved true. Old English Game Bantams have come to stay, and at the present time there can be no gainsaying the fact that they are one of the most popular breeds in our midst. One has only to take a glance at the entries at a few of the leading shows to prove this. The improvement effected in the breed during recent years has been wonderful. The many almost perfect specimens to be seen at such shows as Kendal, Cockermouth, Wigton, Cleator Moor, and some of the South Wales, Midland, and Durham Shows are a treat to behold, and this speaks well for the skill and enthusiasm of the breeders of this popular breed.

Of course, like the Old English Game Fowl, there is no stint in colour or variety, and therefore no lack of choice for anyone taking up the breed. Spangles hold pride of place as the popular colour, and this variety appears to

increase its popularity as time goes on. Black-reds come next in favour, though in recent years they have been closely pressed by some of the off-coloured varieties. The varieties most commonly seen are Spangles, Black-reds, Blacks, Blues, Blue-reds, Duckwings, Birchens, Brown-reds, Furnaces, Brassy-backs, Creles, and Piles, both plain and muffed, and though these do not comprise the whole of the colours of this sturdy little breed, they are sufficient for the work in hand. Any other information can always be gleaned through the query column of *The Feathered World*.

The breeding of Old English Game Bantams is very interesting, as, owing to the variety of colours, the matings are very extensive, and especially in what may be termed the off-colours. In formation the birds are very different from the moderns, being cobbier in build, and lower on the leg. Broad fronts, well placed shoulders, and shape generally should be big factors in choosing birds for breeding. Taking Black-reds first, these can be bred from Partridge or Wheaten hens, though the majority of breeders keep to the Wheaten strain. Partridge breeding, however, is increasing, and that fact accounts for classes being put on for this colour. It is not wise to mix the two colours, for though the cockerels would come all right, the pullets would be practically useless. The soundest and brightest coloured cocks are those bred from Partridges, but are perhaps more inclined to be a bit striped in hackle. It will be seen, therefore, that it is best to mate a Partridge-bred cock to Partridge hens, and a Wheaten-bred cock to Wheaten hens. Both breed fairly true to colour. Brightness of top colour and soundness of black are necessary in cockerels. Partridge pullets should be free from coarseness of pencilling in plumage, of a nice level shade of colour, and clear of rustiness on wings, as far as exhibition points are concerned, though the rusty-winged ones are all right for cockerel breeding.

In Spangles, the rich dark colour is the one most favoured, and these can be bred from Spangles on both sides, or from a Black-red cross. In breeding Spangles, I think if even-coloured birds are selected on both sides, the produce both in cockerels and pullets will be found satisfactory, as, unlike many colours, one pen will throw good specimens of both sexes. Should, however, the chicks come too light in colour it will be found of advantage to introduce a Partridge hen into the pen.

Blue-reds, or Blue-duns as they are sometimes called, are perhaps not seen as much as they deserve to be. They can be bred from Blue-reds, or from a cross with the Black-red, and also from one or two other matings. Duckwings are bred from pure Duckwings, and also from a Black-red cross, and occasionally from Blues, thereby getting what are known as Blue Duckwings. Blacks can, of course, be bred from Blacks on both sides, and richness and soundness of colour should be looked for in the stock birds. In the breeding of Blues,



OLD ENGLISH GAME SPANGLE BANTAM COCK.

The property of Messrs. Read & Newton.

Blue and Black are often intermingled, i.e., mating a Black cock to Blue hens, or having a Black hen amongst the Blues. This is necessary to retain a good depth of colour. In the breeding of Piles, Birchens, and Brown-reds, a similar line of procedure can be adopted as advocated in the breeding of moderns. Creles can also be bred from Creles on both sides, though probably these would throw a small proportion of mismarked ones. Still, this to a big extent would depend entirely as to how long they had been pure Crele-bred. A Partridge hen can be mated to a Crele cock, and this mating produces some excellent Creles. Blue-red and Self-blue hens are also used in breeding Creles, and these tend to give a little

more lustre to the colour. In the off-colours, many matings can be gone in for, and as long as shape, type, and size are secured, colour is not of such great importance.

The standard as drawn up by the Old English Game Club is as follows :—

Head, strong, bold, medium length; eyes, large, bright and prominent, full of expression, and alike in colour; beak, strong at base, and slightly curved; comb, small, both in cock and hen, and evenly serrated at its edge, single, erect, and of fine texture; wattles, fine texture, and small; ear-lobes, to match the comb and wattles as nearly as possible; face, of fine texture, to match the comb and wattles; neck, long and very strong at junction with body; neck hackle, wiry, long feathers, covering the shoulders; breast, back, and body, short back, broad across the shoulders, tapering well to the tail, with a full, broad, well-rounded chest, showing as little keel as possible; the keel or breast-bone to be straight and of medium depth, tapering well up behind, giving a very small and compact belly; the whole body with wings as seen from the top to appear flat and as near heater shape as possible; wings, full and round, inclining to meet under the tail, with strong prominent butts, feathers to be fairly broad and furnished with hard strong quills; the primaries not to be too long, and to be nicely rounded at the ends and to

project past the body as little as possible; Ro tail, in the cock, to be carried at a nice angle, for neither too low nor too high and to be straight; the wry or squirrel tails to be considered a serious defect; feathers to be broad and strong, with Ba a pair of good curved sickles of fair length, n and well furnished with side hangers; in the ed hen, well carried, fairly close, and of medium th length; legs, thighs short, thick, and muscular, wh and set fairly well apart; shanks of medium th length, with good round bone, not flat on shins; in in-kneed or bow-legged to be considered a serious fault; feet, four toes on each foot; should be clean, even, long and spreading, the back toe standing well backward and flat on the ground; spur, low on the leg; carriage, bold, smart, the movement quick and graceful, proud and sprightly, as if ready for any emergency; handling, flesh firm but corky, with plenty of muscle, this to be considered an important factor; plumage, hard, glossy, and firm.

SCALE OF POINTS

Shape and carriage	..	..	40
Handling and condition	..	..	15
Head and eye	..	..	10
Legs and feet	..	..	15
Colour and plumage	..	..	20
			100

Standards can be had of the Secretary of the Club.

CHAPTER III

VARIETY BANTAMS

ROSECOMBS

HAVING gone fully into all the principal breeds of Game Bantams, I will now go on to the Variety Bantams, which are quite as popular as the Game, and as a rule present less difficulties to the young fancier in rearing and preparing for exhibition, with the exception of Rosecombs, which are undoubtedly the most popular of Variety Bantams.

At the present day, competition in Rosecombs is excessively keen at all shows, and a bird to be successful at any show must approach very nearly to the ideal standard of perfection, or his chance of success in the prize list will be a very poor one. There is no doubt whatever but that The Variety Bantam Club and The ye



PARTRIDGE O.E. GAME BANTAM.

Bred by Dr. Perry-Walker.

; Rosecomb Bantam Club are largely responsible for the rapid strides this breed has made during the past few years, as well as doing much good work in furthering the interest of all Variety Bantams. Yet there is, I am sorry to say, much more to do, and although I shall undoubtedly offend some of my readers, I candidly think there is no other breed except Hamburgs, which is improved by science and art more than the Rosecomb Bantam. I do not hesitate in saying that quite 40 to 50 per cent. of the

Black Rosecombs are fairly hardy, but the chickens require very careful attention during the first few weeks, as they are at times difficult to rear, especially so on a cold, damp soil. They are capital layers, and not very apt to become broody.

I will now describe what is considered an ideal Rosecomb cock: Comb fitting close to head, fairly broad and full in front, tapering to a long leader or spike at the back, the leader to be fine, and have an upward tendency



Photo]

BLACK ROSECOMB BANTAM COCK.

One of Lord Dewar's winning midgets.

[Sport and General.

present-day Rosecombs are manufactured, not bred. That is a large percentage, but it is nevertheless correct, and yet these birds go on winning and will do so until either The Variety Bantam Club or The Poultry Club put their foot down firmly in the matter. It seems a great pity that it should be so, for there is no more handsome bird on the whole poultry list than Black Rosecombs, which have been before the Fancy for nearly fifty years.

towards the end; the comb to be well and evenly serrated, not hollow in front or centre, but level throughout; face cherry red, smooth, and free from white, which is often found just under the eye; head short and fairly broad, beak dark and slightly curved; eyes dark and full, ear-lobes white, large, and round, of smooth texture, and the thicker the better, although a bird with a large thick lobe is liable to go white in face as a rule; wattles round and smooth, neck short and thick,



BLACK ROSECOMB BANTAM PULLET.

The property of Mr. Edgar P. Morton.

full hackle, covering the shoulders well; back short and broad, wings fairly large, but not too long; tail large, primary feathers long and broad, sickle feathers long, and broad as possible, the end of the sickle to be almost as broad as the commencement, not running to a fine point. The sickle should be gracefully curved till it almost touches the ground. The side hangers which cover the tail should correspond with the sickle, as broad as possible, and the more of them the better. The breast should be broad and full, and carried forward, with the head thrown back. Legs short and fine, black in colour, not slaty-blue or pale, although adult birds will be found paler in legs than cockerels, and either black or white toe nails, latter preferred. Carriage jaunty and lively, yet graceful in movement; weight, 14 to 18 ozs.; colour lustrous, greenish black, the brighter the better.

Hen: Comb, head, face, eye and lobes same as described in cock; ear-lobes of nice size, the larger the better, if smooth, white, and of firm texture; wattles fine, round, and bright healthy red in colour; back as short as possible, tail carried rather gay, wings carried slightly below the body, not too long; breast full and carried forward as in cock; legs short and fine and black in colour; carriage jaunty yet graceful.

Having described my ideal I will now try and instruct my readers how to mate up to obtain these requirements. Before doing so, just a word of advice. If you contemplate taking up this variety, go in for some reliable

strain, and keep that strain pure. Do not buy from first one and then another, and hope for success; if you do you will regret it, for the only sure road to success in Rosecomb breeding is by keeping one strain pure. Of course it would not always do to breed year after year from your own, unless you retain a certain number of stock birds annually for this purpose, which is done by our most successful breeders, and then this must be done judiciously. Let the health of your chickens guide you as to the necessity of fresh blood in your stock, and then it is advisable to use only those with a dash of your own strain in them, rather than a direct cross, as by using quite a new strain you will in all probability get your birds much too large.

To breed exhibition Black Rosecomb cocks, I would select a cock with a long comb nicely worked throughout and tapering off to a long leader, lobe of good size, round and of good texture; body small, and short in back; plenty of feather, and low on leg, which should be black. To this bird I would mate three or four hens or pullets, former preferred, all possessing good combs and long in leaders; this is important. The lobes should be thick and smooth, and of nice size; face free from white, and dark in eye and dark legs. Body as short as possible, but plenty of tail, and see that the tail feathers are broad, and if the top feathers on each side of the tail are curved all the better. From this pen you should get a good percentage of first-class exhibition



BLACK ROSECOMB BANTAM COCK.

The property of Mr. R. Fletcher Hearnshaw.



WHITE ROSECOMB COCKEREL.

Exhibited by Mr. W. Bradley.

cockerels. Although it is quite possible to breed both good cockerels and pullets from one pen, still, it is advisable, where practicable, to use separate pens; for to breed from one pen you must have a thorough knowledge of how each individual bird is bred and whether of a cockerel strain or a pullet strain.

In pullet breeding the male bird should have a big lobe, as thick and as smooth as possible, *but* must be free from white in face, both these requirements being of great importance. He must also be short and cobby, and by no means long in back. The hens to mate with him should approach as near as possible to the ideal already described. The produce from each pen should be marked and kept separately for future breeding. From the pullet pen, should you by chance breed a cockerel good in head points and shape, yet faulty in colour, viz. red or bronze in saddle or back, don't go and wring his neck as a "waster"; if so, you are killing the goose that lays the golden egg, as it were, for it is these bronzed-backed, red-hackled ones which breed the richest coloured birds. You didn't know that before, did you? yet it is a fact all the same. Here endeth the Blacks.

Whites, like the Irishman's promotion, have had a downward tendency the last few years, so that at the present time a really first-class White is very rare indeed. This variety appears

to have been greatly neglected, but why, I am at a loss to tell, as it is a very pretty and taking variety. True it requires more care and attention in preparing for the show-pen and preserving its colour, which is liable to become tanned by exposure to the sun and weather, still, where kept under favourable circumstances, there is not a more taking variety. The points required are the same as in Blacks, with the exception of the colour, which should be pure white, not yellow or creamy on back; legs, beak, and feet should also be white; lobes white; face, comb and wattles, bright red.

In mating up Whites the same lines must be followed as in Blacks, and where lobe and length of feather have been deficient, these have been remedied by crossing White hens with a Black cock, and for this purpose it is best to select a male bird that is inclined to be slate coloured in leg, the paler the better, but possessing plenty of feather and large lobe. The produce from this pen would come all self-colours, i.e. Blacks or Whites, not mottled; the Whites would in all probability show a little duskiness in leg colour, but this could be remedied the following year if bred back, viz. the White hens mated to the best of the cockerels, and the pullets mated to a White cock; by doing this you would greatly improve the lobe and length of feather.

SCALE OF POINTS

Comb	..	..	..	..	20
Ear-lobes	..	..	..	..	15
Tail ..	..	..	..	..	15
Type	..	..	..	..	15
Condition	..	..	..	..	15
Colour	..	..	..	..	12
Legs	..	..	..	..	8
					100




CHAPTER IV

COCHINS OR PEKINS

THERE is no doubt whatever that next to the Rosecombs in popularity comes the Pekin, or what was formerly called the Cochin China Bantam, owing to the variety having originated from China, the first pair, we believe, having been imported from Pekin about fifty years ago. At that time Buffs were the only colour established, but to-day we have Buffs, Blacks, Whites, and Partridge; to say nothing of Mottles or Ancona Pekins, which are the very latest additions to the breed. The original Buffs were quite different to the present-day type, both in shape and colour, being higher

this is because they are the more easily got ready for show, very little washing being necessary as in most of the other colours. Whites appear to be declining, though during the past season a little more life and enthusiasm have been manifested than have been seen for a long time. Buffs, too, have been on the downward grade, though during the last two or three seasons there has been a determined attempt on the part of breeders to put this colour on the high plane it once occupied, and their efforts have been eminently successful. Cuckoos and Mottles are now very rarely, if ever, seen, and it is only occasionally that one gets a glimpse of a Partridge. These three varieties have never been very




BLACK COCHIN OR PEKIN COCK AND HEN.
Both bred and exhibited by Messrs. Wright Bros.

on leg, longer in back, and more inclined to be what is termed "vulture hocked," more after the style of the booted Bantam. Through the introduction of the Cochin blood by breeding down from the larger breeds, the shape and feather, as well as colour, have been greatly improved, whilst the size also has been reduced.

With the exception, perhaps, of the very best exhibition specimens of Buff Cochins, the Pekin Bantam can lay claim to superiority over all the other varieties of Cochins, from an ideal standard of perfection point of view. Blacks have for a long time been far the more popular of the Pekin tribe, and at the present time I do not think I should be exaggerating to say that more Blacks are bred than of all the other colours put together. Probably

popular, and it is hardly likely they will ever become so.

Pekins can be kept under almost any circumstances, if they are liberally supplied with fine sand for the floor of their houses and runs to protect the foot-feather, which is the only drawback to the variety. They are capital layers of tinted eggs, they lay early in the season, and the chickens are easily reared if the breeding stock are healthy. Of course, like all heavily-feathered birds, unless the foot-feather of the male stock bird is cut short, there will be a large percentage of clear eggs, but this can always be remedied by cutting the foot-feather short in cocks and round the vent of the hens, and for this reason it is best to keep birds, especially for stock, apart from the exhibition ones. This is done by some




WINNING VARIETY BANTAMS.

The property of Mr. H. Hough Watson.

of our most successful Cochin breeders, and is a system that I can recommend to everyone, as by doing so you will not only have a greater percentage of fertile eggs, but the chickens will also be much stronger and a pleasure to rear.

Pekin hens are capital sitters and mothers; they very rarely lay more than a dozen eggs before becoming broody—this is one of the drawbacks to keeping Pekins. Again, for breeding the very best broody hens by crossing the Pekin hen with a Silky cock, their produce cannot be equalled. There is little wonder then that Pekins are such favourites, as even the "culls" in pullets will fetch good prices as brooders, and find a ready sale at this through an advertisement in *The Feathered World* during the hatching season, from January to June. Sitting hens suitable for Bantam eggs appear to be very scarce, and I am surprised that someone does not go in for these as a speciality. Where the fancier has plenty of accommodation I feel sure it would pay well. Unlike the Rosecomb mentioned in my last, the Pekin offers little scope for the habitual faker or trimmer, apart from its tail, which, I am sorry to say, is more often left at home than sent with the bird to the show. Some years ago this practice engaged the attention of The Poultry Club, and this body passed a rule that the birds of this breed should be shown with full tails in. Since then the habit has, to a big extent, been broken off; still, there is room for just a little improvement.



BLACK PEKIN BANTAM.

Bred and exhibited by Messrs. Hallett Bros.

A Game Bantam or a Rosecomb Bantam with a pulled tail would not be acknowledged by the judge; then, why should the Pekin be an exception, and be allowed in the prize list, as 50 per cent. of the present-day Pekins are?

I will now describe what I consider an ideal Pekin, and, for simplicity, give the shape and type first, which are the same in all the different varieties. In the first place, the Pekin is intended to represent the large Cochin in miniature, and should therefore be as near as possible the same shape as the Cochin. Commencing at the head, the comb should be single and finely serrated, and the smaller the comb the better, although it is almost an impossibility to breed Pekins as small in comb in comparison to the Cochin: eyes red, or nearly red, face red and smooth, lobes free from white, neck short and thick, chest broad and carried well forward, back short and broad, the shorter the better; cushion full and nicely rounded, tail short and full, the feathers of the tail proper, twelve in number, being much softer in the quill than in any other breed of Bantam, and should be curved not straight, as in other varieties. In addition, the cock's tail should be well furnished with side hangers corresponding in colour with the body. The wings should be short, and tightly tucked up, legs short as possible, so that the body almost touches the ground; the legs should be abundantly feathered right to the end of the middle toe without being "vulture hocked," which is a great defect in all Pekins. The colour of legs in all the varieties of Pekins should be yellow, and the richer the yellow the better, although it is a difficult point to get in Blacks. Willow or green legs should count heavily against the bird in the show-pen. The one prevailing weakness in all Pekins is scaly leg, but this can be prevented by proper care and attention in keeping the houses and runs dry and clean. Where the birds are kept on nice dry sand scaly leg will rarely appear.

Colour will now engage our attention, commencing with *Bufs*. The exhibition cock should be a rich dark orange, not red; one uniform shade throughout, including tail and flight feathers, when the wing is opened out. The fluff should also be solid buff, not white or dusky, which is often the case. White or black in the flight feathers or tail is a serious defect, more especially white, and a cock showing this defect should never be retained as a stock bird, for in all probability a great



DUCKWING.

PILE.

BLACK-RED.

BROWN-RED.

MODERN GAME BANTAMS.

percentage of the produce would be useless. The hen to match the cock should be a rich golden buff, just a shade lighter than the cock; or, to be more definite, should correspond as near as possible with the breast colour of the cock. This is also a capital guide in mating up the breeding-pen. The colour should be even throughout, not paler in breast than in body colour, but one rich level buff all through, free from white or black, and buff in under-colour or fluff as well.

From outward appearance a bird may seem a solid buff, but when examined carefully may be found slaty in fluff, sometimes almost black. The flight feathers when the wing is opened out may contain more black or white than buff, and the tail likewise. It is, therefore, really necessary in judging Buffs to examine each specimen carefully for these defects, which are often hidden by outward appearances. In mating up Buffs, preference should be given to a male bird of the darker shade, providing he is pure in flights and tail, especially free from white, which is highly important. When I say darker shade, I don't mean a red or dark cinnamon, but a rich deep orange buff of a level colour, not dark in saddle and pale in breast, the richer or deeper in under-colour or fluff the richer will be the colour of the produce: this is one of the most important features in Buff breeding. The bird may appear to be an excellent top colour, but on examination his under-colour may be white or nearly black, both of which are detrimental to producing satisfactory offspring.

The hens to match this cock should be as rich and level in colour as possible; don't breed from pale mealy hens, or those showing a mixture of white in the feather, and, above all, avoid those white in flights or tail. By selecting a pen as described, say not more than

four hens and a cock, exhibition specimens of both sexes can be produced from the same pen. Unlike most varieties of Bantams, the Pekin does not attain perfection until after the second year, as it is not fully developed until twelve months old. They can be hatched as late as July with success, and by hatching late the size and length of leg are reduced, both important features in Pekins.



WHITE COCHIN, OR PEKIN, BANTAM COCK.

Bred and exhibited by Mr. H. Hough Watson.

Blacks should be the same as Buffs in comb, face, eyes, and legs. The plumage should be a lustrous beetle-green as seen in the Black Hamburg, although it is hardly possible to obtain the same sheen as in Hamburgs, but the more lustre the better, as colour in Blacks is a very important point. The fluff of under-colour should also be black to the skin, not grey, which is often found in Blacks, although

birds with sound black fluff are few and far between, still, if bred pure and not allowed to cross with the Whites, they can be bred sound in fluff; it is the crossing with Whites to strengthen the strain and obtain the length of feather that is the cause of grey in under-colour and hackle, which we frequently find in Blacks. To breed Blacks, although it is quite possible to breed exhibition specimens of both sexes from one pen, I should advise, where practicable, to mate up a cockerel-breeding pen separate from the pullet pen. In the cock pen I should select the shortest backed bird I could find, one possessing plenty of feather on shanks and around the hocks and carried out to the end of the centre toe. I should like him low on the leg, excelling in lustre and free from white or grey in under-colour and foot-feather. The male bird in the pullet pen should be descended from a pullet-breeding strain. A good guide in the selection of cockerels for pullet breeding is to notice if he possesses, more or less, a tinge of red in the saddle and neck. If this is found, you can mate him up with confidence that

he will produce the glossy sheen in the pullets, but the hens in both pens should be free from any foreign colour and as shapely as possible, discarding those grey or white in fluff or white in lobe.

White Pekins have found many admirers during the past few years, and are making steady progress. When kept under favourable circumstances—i.e. shaded by covered runs, or kept in small woods or orchards from the sun's rays—White Pekins are a very pretty variety to keep, but should never be attempted where the only accommodation is a backyard. To be successful with Whites they must have grass and a run with covered top and a portion of the sides and front, whilst the floor of the house should have fine sea-sand, to the depth of three or four inches, to protect the foot-feather and keep it clean. Whites are easy to breed true to colour by keeping them pure and in-breeding, but should your chicks come weakly through too much in-breeding then the best plan would be to introduce a sound coloured Black cock, and mate him to the White hens. From these you would get both Blacks and Whites; the former would be more or less grey in under-colour, and show perhaps a little white in foot-feather, which, as a rule, disappears before the bird is exhibited; the Whites will be found very pure in colour, although perhaps not quite so rich a yellow in leg colour, but the produce will be found stronger than the previous year, and more feathers will have been obtained as well. It would not be wise to breed from the Blacks produced in this way, even should they be perfectly sound in colour.

Cuckoo Pekins are a much later addition to the Fancy than Blacks or Whites, and do not appear to make any headway at all. Doubtless this is greatly due to the difficulties of producing them true to colour. It is very rarely we find a single first-class exhibition specimen at any of the provincial shows. In markings they should be similar to the Scots Grey Bantam, the ground colour should be a very pale blue, with dark (almost black) bars; the more clearly defined in barring (as in Plymouth Rocks) the better. Like the Scots Grey and Plymouth Rock, the ground colour varies considerably, but the colour should be as uniform as possible, and the barring fine and distinct; and not only should the top feather be barred, but the under-colour and foot-feather as well. This is very important when mating up the bird for breeding purposes.



Photo]

[Hosegood.

WHITE COCHIN BANTAM COCK.

Bred and Exhibited by Mr. J. F. Entwisle.


I have seen perfect top-coloured birds entirely devoid of fluff barring; but these are not the birds that should be selected for the breeding-pen, and, again, never select birds that show any great amount of white in tail or flights. This is a great weakness in all barred varieties, and should be strictly guarded against.

In mating up the breeding-pens, it would be as well to use two pens. For the cockerel-breeding pen, select a very uniform coloured bird of the medium shade. See that his tail and sickle feathers are well barred down to the root, and that his flights when opened out are sound and free from white. With him mate up three hens or pullets of a nice level colour and clear in ground colour. In the pullet-pen I should use a lighter shade of colour in the cock, but not brassy on top. This is a serious fault. See that his breast is clear and fine in markings, not blurred or indistinct; a little dark on tail is no objection, but he must show no trace of white either in tail or wing. By mating him to three or four distinctly barred hens, nice and clear in ground colour, and especially sound in the black of the barring, you should be rewarded by a good percentage of exhibition pullets, and the cockerels the following year would be found useful if mated back to the hens again, as well as breeding cockerels and pullets together. By doing this you will have two strings to your bow, as it were. Should any of the pullets come black, by mating these to a finely barred cockerel of the lighter shade you would in all probability breed the very best of Cuckoo pullets, and pullets bred in this manner are very useful as future pullet breeders.

We have now come to *Partridge Pekins*, which are intended to represent the Partridge Cochins in miniature, though as yet they have made but slow progress towards the ideal, and have not made many friends. This is no doubt owing to the many difficulties experienced by those who have endeavoured to establish the correct colour and markings in the pullets, combined with length of foot-feather and small size, all difficult points to overcome, although with care in mating and in breeding there is no reason why the desired requirements should not be obtained, although it will require a lot of patience and perseverance, both essential traits in every would-be successful fancier's life, and without which no fancier

can hope to reach the top rung of the ladder of fame in the Fancy. To describe the colours of the Partridge Pekin cock I cannot do better than refer my readers to the top colour of the Black-red Game Bantam cock. Both these birds are identical in colour, except in hackle, where the Partridge Pekin shows more striping and is generally two shades of colour in hackle, dark orange towards the head, shading off towards the shoulders to pale golden. The breast and thighs should be a sound black of a greenish shade, legs and feet yellow, and well furnished with black feathers right to the end of the centre toe, and as free from white foot feathering as possible, and the longer the feather the better. The hen should be a rich chocolate ground colour, of one even shade throughout, every feather evenly pencilled with black, except hackles, which should be a light golden, the ground colour and pencilling being two of the most important and difficult points to obtain.

To breed Partridge Pekins it is desirable to have two pens, as it is next to impossible to breed exhibition cockerels and pullets from the same pen. In the cock pen you would require



PEKIN BANTAMS.

Bred and exhibited by Mr. J. S. Kirkland.

to have the male bird particularly bright in his top colour, and sound black in breast, thighs, fluff, and foot feather, otherwise your cockerels would be rusty in fluff and breast, and white or grizzly in foot feather. The hens to mate with this cock should be a pale in hackle as possible; they need not be particularly good in pencilling, as this is not necessary in cockerel breeding, but see that they are well feathered, and especially so on middle toe. Now, to produce the best exhibition pullets you should select a cock a shade darker than the above, and one that is coarser in his striping of hackle; in all probability he will be more or less faulty in breast colour and fluff, showing red or rusty-coloured feathers, if so all the better for the purpose you require and a capital guide that he is a pullet breeder. To him mate three or four of the very soundest, ground-coloured hens you can find, well and distinctly pencilled, and from these you may expect a good percentage of sound coloured pullets.

In mating up both pens always have the proper shape in your eye—viz. broad in chest, short in back and leg, low on the ground and abundantly feathered, all necessary points to be considered when mating up.

In the first place strive for colour and type, and when you have mastered these requirements you can always reduce the size and retain the colour, etc., by inbreeding.

SCALE OF POINTS—COCHINS OR PEKINS.


Colour	20
Type and carriage.. ..	15
Fluff and cushion	15
Head	10
Leg and foot feather	10
Leg, length.. ..	10
Condition	10
Size and weight	10

100



CHAPTER V

BRAHMA BANTAMS



Photo] "Feathered World."

DARK BRAHMA BANTAM
COCKEREL.

The property of Mr. J. H. Firth.

THIS is another variety of the feather-legged tribe which has been before the Fancy for the past twenty years or more, but only in the last few years have they made any big strides, the credit of which must be put down to the club formed for the breed. Classes are now provided at most of the leading fixtures, and the entries compare favourably with the majority of

other breeds. Although a very pretty and hardy variety, I do not think they will ever become as popular as either the Pekin or Booted, principally on account of the difficulty of breeding them true to colour, and at the same time keeping down the size and producing the required length of foot feather, a point in which fifty per cent. of the Brahma Bantams of to-day fail.

Like the large Brahmas, there are two varieties only, viz. lights and darks, the former as yet being most favoured, doubtless on account of being the easier of the two to breed true to colour. Still, I consider the darks equally as handsome, in fact a very taking variety.

The hens are good layers of a tinted egg, and the chicks are fairly hardy, and easy to rear.

In shape they should resemble the Pekin as near as possible, except that the Brahma is a trifle higher on the leg, and the hens especially being longer in back and tail. The light cock should have a triple or pea comb; face and wattle, red; hackles, silvery white, striped with black towards the bottom; the back, wings, shoulders, breast, and thighs, white; tail, black; the top outer feathers, or what are termed hangers, slightly edged with white; legs, yellow, heavily feathered with white

feathers to end of middle toe, the more free from black the better; beak, yellow, to match the legs; eyes, red or yellow.

The hen is white in body, with neck hackle also white, but plainly striped with black; the wings when opened out should show black in primaries, and also in under-secondaries; tail also black, to match the cock; legs, feet, and beak, yellow, and well furnished with white feathers to end of toes; eyes, red or yellow.

The dark Brahma cock should have a triple or pea comb, face red, also wattles and lobes, latter free from white; beak and legs, yellow or horn colour; neck hackle, silvery white, striped with black, the stripe to be broader towards the bottom of hackle; breast, thighs leg and foot feathering, tail, wing-butts, and shoulders, sound black; wing-bow, back, and shoulders, silvery white; the saddle hackle same colour, but with distinct black striping; wing-bar and tail-hangers, rich green-black; the leg

and foot feathering to be a sound black and as long as possible—the less white in foot-feather the better, although it is a difficult matter to get the foot-feather absolutely sound in either colour, as black will invariably show in the foot-feather of the lights, but it should always be guarded against when mating up the breeding-pen. This only refers to the cocks, as the hens are generally fairly sound in this respect. The dark hens should be identically the same as the cock in face, eyes, and leg. Hackle, silvery-white ground-colour distinctly striped with black, tail also black. Body, breast and wing colour to be a nice steel-grey colour, pencilled with black, the pencilling to be as distinct and as even as possible and continued right up to the throat and out towards the fluff behind the legs; the pencilling and ground colour are the chief and most important parts in dark Brahmas, together with length of leg feather, which should be as long as possible and carried right to end of middle toe.

And now to the breeding of Brahma Bantams. Although the proper ground-colour of the dark hens is hardly properly established yet, it would be advisable to mate up two pens to obtain the best results. To breed the best cockerels it is always best to have the soundest coloured bird you can possibly find; one sound in his black, and clear in his top colour. See that he is well feathered and good in shape, but not large, the smaller the better, as size is as yet a failing point in Brahmas. With him mate three of the best-shaped pullets you have, good in head points and comb, and the lighter and more silvery in hackle to top of head the better for cockerel breeding; although their body and breast colour may be very pale and defective in lacing, this is no detriment in cock breeding, so long as the neck hackle is a very pale silvery-white.

In breeding pullets the order must be reversed. See that your hens or pullets are sound and uniform in ground colour, also regular and even in pencilling. This is absolutely necessary. The cock should be as described before, except that if he shows a little pencilling on breast and fluff, and also wing bar, all the better for pullet breeding; the more lacing he shows proves him to be descended from a pullet breeding strain. From this pen it would be advisable to retain three or four of the best and soundest-coloured pullets, also a couple of the best-shaped cockerels, and the following



Photo by]

[J. Drew, Aldershot.

LIGHT BRAHMA BANTAM PULLET.

The property of Mr. J. Greaves.

season breed from these. Then you would be able to establish a thoroughly reliable pullet-breeding strain, and at the same time you are reducing the size and improving the colour.

To breed lights it would scarcely be necessary to use two pens, as both cockerels and pullets might be produced from one if properly mated. Take a cock that is as near perfect in colour as possible, not too heavily striped in hackle or too pale, but use one of the medium shade, but see that his striping in hackle is sound, rich black, not tinged with grey, and finely-striped saddle hackle. He must have plenty of feather, and be a good shape, with a small, neat comb. With him mate a couple of hens or

Brahma Bantams require the same housing and attention to the protection of the foot feather as described in last article on Booted, and may be bred to advantage until May, June, or July.

SCALE OF POINTS.

Head and comb	..	..	..	10
Length of feather and fluff	..	..	..	20
Colour	..	..	..	20
Shape	..	..	..	20
Size	..	..	..	15
Condition	..	..	..	15

100



[Photo]

LIGHT BRAHMA BANTAM COCK.

[Gunston, Braajord.

Bred and exhibited by Mr. J. Greaves.

pullets as clean white on back and wings as possible, and if a little fine in striping or hackle all the better; these are for breeding cockerels. Also two hens that must be perfectly sound on back, a pure white, and the whiter the fluff or under-colour the better; hackle well striped with black, the broader and more distinct the striping the better; these for pullet breeding.

CHAPTER VI

MALAYS

THE Malay Bantam is undoubtedly descended from the large breed of Malay, and during the past few years has greatly improved in type, *but there is yet much room for further improvement*, for as yet the majority of the present-day

winners are much *too large*. If they could only be reduced in size, and at the same time retain the true Malay type, I feel certain they would soon find many new admirers. In my own mind I am firmly convinced that some Malay breeders have made a great mistake by introducing the modern Black-red Bantam in order to reduce the size, but by doing so they have lost the type and carriage peculiar to the Malay. My contention is that in judging Malay Bantams the first things to look for are type, shape, and carriage, then head points, size, and colour, for the latter is only what one might call a secondary consideration, whilst type and carriage are absolutely necessary. No matter how good the bird may be in colour and size, if it lacks the general character, type, and carriage of a Malay fowl, it should not be countenanced in the show-pen.

Malay Bantams are generally much too large, although there are here and there small and typical specimens to be found, at our big shows in particular. If fanciers would only follow a system of breeding by first getting the proper type and general characteristics, irrespective of size, and afterwards inbreeding from these typical specimens, the desired size as well as type will be obtained in the long run, and much quicker than by introducing Modern Game Bantams, which can never have the desired effect, and bound sooner or later to end in complete failure.


There are four varieties of Malays—Reds, Whites, Piles, and Blacks, although it is very seldom we find either of the two last named at any of the shows. The Reds and Whites have found many friends, and the winners are widely spread, which is one great advantage to the welfare and future prosperity of the breed. The Malay Bantam should be an exact miniature of the Malay fowl.

The head should be broad, with overhanging eyebrows, giving the bird a fierce expression. The beak should be strong and curved, horn or bright yellow, the latter preferred. Eyes yellow or daw, and well sunk in the head. Comb small, walnut shape, and set well forward on head. Face red, smooth, and free from feathers. Neck as long as possible, and carried very erect. Hackle short and fine. Back fairly long, and sloping towards the tail. Shoulders broad, and carried high and well forward, having the appearance of being apart from the body, the shoulder points to be very prominent,

which gives the bird a bold appearance. Wings short and strong, and fitting closely to the body. The tail of the male bird should be drooping, medium length, narrow in feather and sickles, which should be slightly curved. The hen's tail should be short and square, and carried slightly above the horizontal line. Breast deep and broad, and almost bare of feathers. Thighs long and powerful, scantily feathered, and set wide apart. Shanks long and fine, round in front, and a rich orange yellow in colour. Toes (four) long and straight, the back toe set on low, and resting firmly on the ground in a direct line with the middle toe. Plumage short, and as hard as possible.

We now come to the colour of Malay Bantams.

There are two distinct shades of the Red, either or both of which are acknowledged in the standard and in the show-pen. The one is much lighter and brighter altogether in top colour of cocks than the other, and should be as near as possible the colour of the Modern Game Bantam which I have described for



WHITE MALAY BANTAM COCKEREL.

The property of Mr. William Arkwright.

pullet breeding, and the hens to mate with this should be Partridge, and if sound in body colour they should produce both good cockerels and pullets, for as a rule they breed very true to colour. The other shade of red is the dark, maroon top-coloured cock. The breast, wing bar, lights, and shoulder-points should be a lustrous black, hackle a dark red, wing-ends or secondaries rich bay. The hens to mate with this shade of bird, which, as a rule, are harder and shorter in feather than the bright reds should be the wheaten or cinnamon of one uniform shade of colour throughout. By mating in this way exhibition specimens of both sizes can be bred from one pen, which is a great advantage to those whose space is limited.

Whites are a very taking variety, their beautiful snow-white plumage and rich orange legs and beaks showing up well together. They are easily bred, providing care is taken not to breed from male birds that are *strawy* or *yellow* on top, for using such birds is simply a waste of time, as the produce will invariably turn out fifty per cent. worse, and as such are useless in the show-pen. In mating up your Whites, then, let your first object be to select a cock pure white on saddle and back, and one free from willow in leg, for this is one of the failings of Whites, but with proper selection in mating up the fault can easily be remedied.

Piles are very few and far between, no doubt owing to the difficulty in obtaining the desired colour in the male birds, but this can be remedied if my instructions in breeding Pile Game Bantams be followed out, viz. take a couple of White hens, and mate them to a rich, deep-coloured Red cock, care being taken that he is a rich dark bay on wing-ends, and particularly sound in his black both on breast and wing bar, and by doing so you will be rewarded by a fair percentage of better Piles than have hitherto graced the show-pen.

The following year I would breed brothers and sisters together, and should the percentage of good birds from the first cross be satisfactory, then by all means go on again on the same lines, as well as by using the best of the Pile cockerels from these, and mating him to his sisters. Never use a Pile cock that is weak in his bays as a stock bird, for it is labour in vain, as they are useless either for stock or exhibition purposes; I would much prefer to use a dark wing-ended bird, even should he be faulty in breast or top, rather than use an otherwise perfect-coloured bird with faulty wing-ends. This reminds me of what a noted Bantam breeder

and exhibitor said to me the other day: "Proud, you are telling young Bantam fanciers too much"; but from the numerous encouraging letters I have received from all parts of the country, I am happy to learn that the practical information I have penned, with a view of giving all young fanciers a helping hand, has been so thoroughly appreciated, that no matter whether I am telling too many of the secrets of breeding, etc., or not, I shall carry out my promise set forth in the opening chapters of this work, viz. that everything that can be of use to the young fancier will be written in language as brief and as simple as possible, that he who runs may read, and may benefit by his reading. It is not that I am telling young fanciers and amateurs too much, but I am telling them, perhaps, something hitherto unknown to them, and which those already in the know would prefer keeping secret.

If you can give a struggling fancier a little advice by the way or put him on the right road to successful breeding after his years of toil, I say do so, and I feel confident that he will be heartily thankful for your assistance; and remember that "good advice costs little, but is worth much."

SCALE OF POINTS.

Type	..	..	..	..	28
Head	..	..	..	..	25
Feather	..	..	..	..	10
Legs	..	..	..	..	10
Gloss and vigour	..	..	..	..	8
Shoulders	..	..	..	..	7
Tail	..	..	..	..	6
Colour	..	..	..	..	6
					100

SERIOUS DEFECTS.—Excessive size, and other defects in connection with Malay fowls.

CHAPTER VII

INDIAN GAME

AMONGST the many varieties of the Bantam Fancy none is more beautiful or more noble-looking a miniature Game fowl than the Indian Game, which, I think, were introduced to the public about the year 1887.

Of course, at that time the variety was much larger than we are accustomed to see now at the big events; and although during the past years



INDIAN GAME BANTAM HEN.

Owned by Mr. L. Ardern.

they have made fairly good headway, there is yet ample room for further improvements both in size and type. There is very little doubt that the Indian Game Bantam owes its origin to some extent to the Aseel and Indian Game cross, and afterwards by in-breeding from the Indian Game late in the season to reduce the size.

Of course, to reduce Indian Game to Bantam size is a considerable undertaking, and means many years of labour, on account of the massive build of the Cornish Indian Game.

Although Cornwall has always been considered the stronghold of the larger breed, still, in Indian Game Bantams, they have not got it all their own way. At most of our large shows classes are provided for them, and which usually fill well; whilst at the smaller provincial shows it is quite a common occurrence to see an Indian Game Bantam figuring prominently in the prize-list in the A.O.V. class.

Now that The Indian Game Club have recognised their merits by admitting them into their ranks, it is more than likely that their future progress towards perfection will be more rapid.


In every respect the Bantam should be an exact miniature of the Indian Game fowl, and as small as possible consistent with type and build.

In judging Indian Game Bantams shape

should be the first essential point, then colour and size; but however good in the two latter points, if the bird does not possess the necessary shape and build it should not be countenanced in the exhibition pen.

Therefore let type be your first aim, then colour, whilst the required size can always be obtained by in-breeding. I will now try and describe what I should consider an ideal bird.

Cock.—Head: Rather long and thick, skull broad and inclined to be a little beetle-browed. Neck: Medium length and slightly arched. Beak: Well curved, strong, giving the bird a powerful expression. Face: Red, fairly smooth and fine in texture, and dotted over with black feathers. Comb: Pea-combed or triple, small and neat, closely set on head, and bright red in colour. Eyes: Full and prominent, pale red preferred, eyebrows slightly arched. General Shape: Thick and compact, back short and flat, broad as possible at the shoulders and tapering towards the tail, but not flat-sided. Breast: Deep and wide but well rounded, and breast bone straight. Wings: Short, clasping the body closely and firmly, carried well forward, high at points, and well tucked in behind. Tail: Medium length, slightly drooping, tail feathers proper, hard and close fitting, with a



INDIAN GAME BANTAM COCK.

The property of Mr. Lawrence Ardern.

few side hangers or tail coverts narrow and glossy. Thighs: Much shorter than Malays, stout, and well-rounded. Shanks: Medium length, strong, and nicely rounded, close-fitting scales, and a deep rich yellow in colour, the richer the better. Feet: Strong, toes straight, long and well spread, back toe carried well on the ground. Plumage: Hard and glossy. General Appearance: Powerful, sprightly, very vigorous, bold, upright carriage, with the back sloping towards the tail, flesh firm and hard in handling.

Colour: Head, neck, breast and thighs black, with a lustrous green sheen; the lower part of the neck hackle a little mixed with bay or chestnut, which, however, should be almost entirely covered by the body feathers; shoulders and wing bows a rich glossy green-black, the shaft of the feather to show chestnut; tail coverts, rich beetle-green; back, green glossy-black broken with rich bay; wing-bar, glossy black; wing-end, bay or chestnut. Defects: Twisted claws, flat shins, crooked breast-bone, rusty hackles, white in hackle, twisted hackle, high on leg, heavy in feather.

The *hen* should correspond with the cock in shape and build, also head points, neck, and legs. Colour: The ground colour should be a rich mahogany brown or chestnut brown, the lacing to be double on each feather if possible, viz. an outer edge of green glossy black, and an inner one enclosing shaft of feather of green glossy black, the chestnut ground colour between the lacings being perfectly clear and free from black or smut. Defects: Same as above.

In breeding Indian Game Bantams it is advisable to mate up two pens, one for cockerel and the other for pullet breeding, as it is next to impossible to produce exhibition specimens of both sexes from one pen.

To breed the best cockerels I should require a typical shaped bird, low in leg, with plenty of substance of thigh, and broad in chest. Let his shoulders be as prominent as possible, and see that he is free from rust in neck hackle and a sound bay on his wing-ends. Never breed from a narrow, flat-sided cock, or 100 per cent. of your chicks will be worthless. The hens or pullets to match the cock (hens for preference) should be a sound, rich dark ground colour, short and dark in hackle, and the more typical in shape the better, and select those short in back and prominent in shoulders; by doing so you may expect the cockerels from this pen to be all that you could desire in colour and shape. The pullets from this pen I would recommend being put back the next season to

the sire and the best-shaped cockerel to the hens; by doing this you are keeping down the size and retaining the proper type and colour as well as breeding up a strain of your own.

In the pullet pen, colour and lacing are both requisite and necessary, therefore select a cock that shows a fair amount of lacing; if he is a bit red in hackle so much the better, but be careful at the same time that you do not overlook the shape and type which are absolutely necessary. In other words, you want a good-shaped, low-legged cock that is faulty in colour i.e. red in hackle and showing lacing, both serious faults in an exhibition cock. With this cock, put about four hens or pullets: it does not matter which in pullet breeding. Select those possessing a rich chestnut ground colour, discarding those pale or washed out in colour; see that the lacing is both double and distinct. Indistinctly laced ones are useless as pullet breeders. The lacing must be a sound rich green glossy-black, hackle must also be sound and free from rust or red. Preference should be given to those low on leg and short in back and the broader in chest the better.

It would be advisable to keep a couple of the best shaped cockerels from this pen which show the most lacing throughout for future pullet breeding. It is much better to build up a strain of your own that can be relied upon, for you can rest assured that if the pullets from



Photo]

[“The Feathered World”]

INDIAN GAME BANTAM COCK.

The property of Lt.-Com. H. G. Nalder.

this pen are satisfactory as exhibition specimens, especially from a colour point of view, you can depend upon the cockerels being all that you would require as pullet breeders, although they are of no use as exhibition birds. Further, it is not wise to introduce a fresh strain, for this would have a tendency to increase the size, and in all probability spoil the colour and type you had already built up. Never use flat-shinned birds. My advice is, build up a strain of your own, and retain three or four of the very best each year for future stock, and if you do so and peg away perseveringly you are bound to come out at the top in the long run, although you cannot reasonably expect to do so the first season.

SCALE OF POINTS.

Shape	..	..	..	..	20
Size ..	..	..	..	..	15
Colour	..	..	..	..	10
Hardness of feather	..	..	..	..	15
Legs and feet	..	..	..	..	10
Eyes	..	..	..	..	5
Head	..	..	..	..	10
Tail ..	..	..	..	..	5
Condition	..	..	..	..	10
					<u>100</u>

Faults to be deducted accordingly.



JUBILEE INDIAN GAME BANTAM COCK.

The property of Mr. H. Hunt.

JUBILEE INDIAN GAME.

Though to many this is a new breed introduced during the past few years, it is over twenty years ago since Mr. H. Hunt, who then introduced it, was first breeding and exhibiting it. It must have failed to take on in those days, for Mr. Hunt let his strain die out. However, the present resuscitation or reviving of it appears of good omen, for there is no doubt but that the breed is going to be a very popular one. There are a good many breeders taking it up, and already a club, fairly strong in numbers, has been formed for furthering the interests of the breed and its adherents.

The general appearance, shape and carriage should be on the lines of the Indian Game. The standard of perfection as regards colour in the cocks is as follows: Head, neck, under fluff, thighs and tail white; breast white, lacing permissible, but no alien blood be introduced; shoulders and wing-bows, white, slightly broken with bay or chestnut in the centre of the feather or shaft; tail coverts, white; back, white, touched on the fine fronds and the ends of the feathers with bay or chestnut. When the wing is closed there is a triangular patch of bay or chestnut formed of the secondaries, which are white on the inner and bay or chestnut on the outer web; primaries pure white except for about $2\frac{1}{2}$ inches of a narrow lacing of light chestnut on the outer web.

Hen: The ground colour is chestnut brown, nut brown, or mahogany brown; head, hackle and throat, white. The pointed hackle that lies under the neck feathers, white; the breast, commencing on the lower part of the throat and expanding into double lacing on the swell of the breast, white, the inner or double lacing being most distinct; the under parts and thighs marked somewhat similarly and running into a mixture of indistinct markings under the vent and swell of the thighs. The feathers of the shoulders and back are somewhat smaller, enlarging towards the tail coverts, and are similarly marked with the double lacing; often in the best specimens there is an additional mark enclosing the base of the shaft of the feathers and running to a point in the second or inner lacing. The tail coverts are seldom as distinct, but with the same style of marking. The primary or flight feathers are white, except on the inner frond or web, which are a little coloured or peppered with a light chestnut; the secondaries are white on the inner web,

while the outer web is in keeping with the general ground colour, and are edged with a delicate lacing of white; the wing coverts which form the bar are laced like those of the body, and often a little peppered. The white lacing mentioned should be as clear as possible, and should look as if embossed or raised.

Defects.—Cock. Crooked backs, beaks, legs, wry or squirrel tail, in-kneed or bent legs, black hackles, flat sides, single and Malay combs, crooked breasts or claws, flat shins, ticked hackles, breast, thigh or tail, bad shape, heavy feathers, smaller of size, long legs and thighs, twisted hackle.

Hen. Similar to those in cocks, also too light, too dark, or mealy ground colour and defective markings.

Weight.—Cocks, 3 lbs.; Cockerels, $2\frac{1}{2}$ lbs.; Hens, $2\frac{1}{2}$ lbs.; Pullets, 2 lbs.

As regards breeding, the points enumerated for Indian Game apply in the main to this breed.

Another variety, the blue-red, has been exhibited, the originator being, I believe, Mr. Fulcher. This colour was exhibited at the Birmingham Show of 1922, but as far as I remember it has not been shown since. The colour is a very pleasing one, but so far it does not seem to have caught on.



JUBILEE INDIAN GAME BANTAM HEN.

The property of Mr. Henry Hunt.

CHAPTER VIII

BOOTED

THESE have never been a very popular variety, and of late years they seem to have lost a lot of ground, so much so that I fancy their breeders can be counted on the fingers of one hand. The Variety Bantam Club and one or two enthusiastic lovers of the breed try to keep it going, but general interest seems to be lacking, and whilst this is so, the Booted is hardly likely to occupy a very exalted position in the Fancy. This is much to be regretted, as a good specimen is a very pretty and interesting bird.

There are several varieties of Booted, but the most popular are Blacks and Whites, but we should prefer the former, owing to the difficulty of keeping Whites pure in colour, especially the foot-feather, which, except under very favourable circumstances, is apt to become stained. In the Blacks it does not show; and, again, the Whites suffer also from exposure to the sun and weather.

Booted Bantams are very taking, and greatly resemble the Pekin. They are fairly hardy and not difficult to rear, the only trouble being to keep down the size. Although of late years this has to some extent been done, still there is room for further improvement in this respect, as we find them too big and leggy as a rule, especially the Blacks. By inbreeding and hatching late in the season this fault could be easily remedied, and, at the same time, the length of foot- and hock-feather could be still retained, a most important feature of Booted Bantams.

Another improvement in Booted which I should like to see is the "carriage," which is becoming far too near that of the Pekin. We do not want Booted to resemble the Pekin at all; instead of the short, close-fitting wings of the Pekin we want to see the long drooping wings carried low, tail elevated, though not too near the head, long, with good length of sickles and side hangers.

Booted are single combed, ear lobes free from white, although white will often be found, but it is a defect and should be guarded against, wattles red, not too large, neck short and curved, back short, wings low and drooping, tail long and carried high, with plenty of side hangers, legs not too long, abundantly feathered, the feathers being much stiffer than those of the Pekin hock feathers as well as footings

to be as long as possible, and carried out to end of middle toe. Carriage more erect than the Pekin. The leg colours of Blacks should be black, and white in Whites.

The mating of Booted Bantams presents no difficulty whatever, and exhibition cockerels and pullets can be bred from the same pen; still in mating up always give preference to birds short in back and leg, and well furnished both in foot feather and hocks—the more feather the better; see that ear lobes are free from white, and comb evenly serrated.

Keep the Blacks pure and give preference to those having some black under-colour down

which would count heavily against the bird in the show-pen.

In addition to the Blacks and Whites we have the White Whiskered or Muffed, which are identically the same as the White aforesaid, except that they are furnished with feathers round the cheeks and under the throat like whiskers; hence the name Whiskered Booted.

There are also a few Splashed or Spangled, but they are few and far between, and do not appear to become favourites, as they are seldom met in the show-pen.

In breeding Booted I should prefer to hatch the chickens from the end of April until the



WHITE BOOTED BANTAMS.

The property of Mr. Kenneth Ward.

to skin, as it is a great mistake to breed from Blacks which are white in fluff or under-colour. On the other hand, should your Whites show a tendency to come yellow or creamy in colour, this can be remedied by introducing the Black for a single season, and although it may produce a bluish tinge in the leg colour, the body colour will be found to be greatly improved; and the following season, if bred back to Whites, the leg colour will again be secured. The Blacks from a cross of Blacks and Whites should never be kept as stock-birds, but should be carefully marked when young to prevent mistakes. By breeding from these Blacks you would find the majority of the chicks another season would be grey in fluff and throwing white in foot, feather and tail, all of which are defects

middle of June, and should inbreed as long as the foot-feather and health of the produce were maintained.

One great drawback the young fancier who takes up Booted will experience is the number of unfertile eggs each season, owing to the length of foot-feather of the male bird preventing him from treading properly. This difficulty is experienced by all heavily-feathered birds, both in the poultry world as well as pigeons.

The only remedy is to keep birds used for stock purposes apart from the exhibition birds, and before putting the cock with the hens in the breeding-pen to cut off the cock's foot-feathers short. This will found to remedy the failure. There is no necessity to interfere with the



SILVER SEBRIGHT HEN.

Bred by Mr. J. M. Philipson.

foot-feather of the hens. Three hens with the male bird will be found quite sufficient.

In keeping Booted, let me impress upon my readers the advisability of having proper houses and runs, for without these success cannot be obtained. In the first place, the runs must be covered at the top, side, and ends, leaving only the front open, and this must be boarded about two feet from the ground. At the bottom of the run I should prefer to have a thick layer of fine seasand, from four to six inches deep. Failing this, get a few sacks of oat husks, or what is known by the name of "seeds," from your miller. Cover the floor of the run and inside the poultry house with these to a good depth, not less than three or four inches. Remove the droppings two or three times a week, and turn the seeds or sand with a rake at the same time. By carrying out these instructions and keeping the floor of the run thoroughly dry you will be repaid by always having the foot-feathers of your birds in perfect condition and your birds clean, which is half the battle in the exhibition pen.

If your birds are put into a damp earth run, or an unevenly boarded run, you can only expect broken foot feather, and broken foot-feather spells failure in the show-pen. Booted do not require perches; they are all the better for roosting on the ground.

SCALE OF POINTS

Length of feather	..	..	20
Colour	..	..	20
Shape and carriage	..	..	15
Size	..	..	15
Head and comb	..	..	10
Leg colour	..	..	10
Condition	..	..	10
			100

Faults to be deducted accordingly.

CHAPTER IX

SEBRIGHTS

THIS variety is one of the oldest and prettiest members of all the Bantam tribe, and has been before the Fancy for a great number of years, having been brought out and perfected by the late Sir John Sebright. There are two colours, Golds and Silvers; the latter are by far the most popular, being easier to breed true to colour. The greatest difficulty in breeding Golds is to obtain the rich bay or golden bay in the ground colour, together with a fine, even, lustrous green black edging to each feather, at the same time keeping the centre of each feather perfectly clear.

The failing in Golds is to run pale in ground colour and coarse in lacing, whilst the latter is very often found to be a dull brownish black instead of a rich green shade of black.

Sebrights are capital layers, but the chickens during the first few weeks are somewhat difficult to rear, except to the professional breeder. Damp and cold are the worst enemies of the Sebright breeder, the chickens requiring a dry shaded run, free from damp, and well protected from the east winds.

I have known some Sebright breeders experience great difficulty in rearing many chicks and having a large percentage of clear eggs, although at the same time I have known others to be just as successful in the same district. What was the reason? Why, the first-named had been doing the round of shows with his stock birds, whilst the latter had seldom or never exhibited the birds he intended to breed from. How often this is the case, and if the young fancier would only bear this in mind, it would save him endless trouble and annoyance during the breeding season. You

cannot burn the candle at both ends. If you want to have a successful breeding season and healthy, vigorous chicks, the less you exhibit your stock-birds the greater will be your success in breeding, not only in Sebright Bantams, but in every variety of poultry.

I will now try and describe what I consider should constitute an ideal Sebright, and will take the *Silvers* first.

In cocks, I like to see the short square back, perfectly flat, breast full and prominent, wings carried low, head small, comb rose shape, fairly broad in front, and tapering off with long fine leader, the latter not to be carried upwards, but should follow the bend of the neck, or helmet shape, as is found in Wyandottes; face in cocks should be inclined to mulberry. Although it is impossible to get the deep mulberry or dark purple in the face of the cocks the same as in pullets, especially the first year, still they should have a certain amount of damson eye-cere, the darker the better. Ear-lobes free from white, but a little blue or purple is an advantage rather than an objection. Eyes dark as possible; beak horn colour, short and stout; wattles fair size and nicely rounded, not folded or creased; legs and feet free from feathers, slaty blue in colour; tail fairly large and well spread, and, unlike the other breeds of Rosecombs, must have no sickles, although it will sometimes be found that some are produced with sickles, and these are invariably the best breeders, and especially useful as pullet breeders, but it is not advisable to use them in the cockerel-pen. The tail should be carried high and the head well thrown back; plumage, pure silvery white ground colour, each feather finely and distinctly laced with rich green black. When I say distinctly laced I mean that the lacing should be even and distinct all round the feather—not narrow round the sides and broad at the bottom, but one uniform width of lacing right round the feather. The tail feathers should be clear silvery white centre, laced as above, but the ground colour of each feather should be *clear*—i.e. free from black, or what is termed “peppery,” and this fault will be found in quite fifty per cent. of *Silvers*. Still, with care and perseverance this can be bred out completely; only it requires a little time and patience. When mating up your breeding-pens clear-tailed birds should always be preferred to those which are “peppered” or “splashed” in tail. The Silver hen is identically the same in colour and shape, the

only difference being the face and comb of the hen should be a very dark mulberry, much darker than the cock; like the cock she should be short in back, small and compact—the prevalent fault in hens is “long backed,” which is a very serious defect, and should be guarded against in the show-pen.

The only difference between *Golds* and *Silvers* is the ground colour, which should be a rich *deep* golden bay or chestnut, not the pale golden we very often find in the present-day exhibits, which are getting much too pale in colour, whilst the lacing, which should be a sound deep greenish black, is more of a brown than a black, and such birds we generally find devoid of lacing on the flight feathers, which failure should count heavily against the bird in the show-pen.

In breeding Sebrights it is much the best to use two pens, one for cockerel breeding and the other for pullets. In the cockerel pen select a sound laced cock, inclined to the heavy lacing rather than fine. He must be particularly good and distinct in breast lacing and tail. This is highly important. The flights or wing-ends of the cock should be



Photo]

[C. Hosegood.

GOLD SEBRIGHT PULLET.

Bred and exhibited by Mr. C. I. Young.

well laced, ground colour of tail clear, and take care that he shows the desired damson colour round the eyes, the more the better. The hens to match him should be lightly or finely laced, but the lacing should be a sound black, not edged with brown. Each flight feather of the wing should be distinctly laced right to the end, tail perfectly clear in the centre, but every feather laced, face dark mulberry.

In breeding for pullets I should prefer the cockerel to be very fine in lacing, but would at the same time require the pullets to be rather heavily laced, as dense a black as is possible to get, and at the same time the lacing must be distinct and even. The cock for pullet breeding should also have a fairly good breast, a good comb essential, clear in centre of tail feathers, each feather laced, and well-laced wing-ends. As I said before, a cockerel with a tendency to have sickles is very useful as a pullet breeder, and the eggs by a "sickled" cock have been found much more fertile than by others. On the other hand, cocks possessing short tails have invariably been found to be poor breeders.

The hens in the pullet pen, in addition to being rather heavily laced, should possess a good amount of lacing down the thighs, tail coverts and tail: this is very important in pullet breeding.

In breeding Sebrights always select a small cock in preference to small hens, as larger hens are always the most productive and the chickens easier to rear than those hatched from eggs from small hens, and on account of the delicate nature of the Sebright chicks it is at all times best to use hens in preference to pullets in the breeding-pen, although if the pullets are a good size and early hatched I see no reason why they should not prove equally as good.

In conclusion, let me remind my readers of the great value of in-breeding in Sebrights, a breed in which colour and markings play such an important part, and which are almost impossible to obtain without in-breeding. If you think by purchasing half a dozen of the very best exhibition specimens from Tom, Dick and Harry, that you are bound to breed the same the following season, you will quickly find out your mistake, for it is very probable that the majority of the produce would be little better than wasters, whereas had you purchased only a couple of good breeding birds both of the same strain I have no hesitation in saying

that the produce of these two birds would be of more value than all those bred from the half-dozen exhibition birds, which would no doubt have cost you four times as much. Therefore start with some reliable strain and keep it pure. As long as your chickens are healthy, you can go on with the same strain; always breed from healthy stock birds only, and when your chickens come weakly and are difficult to rear, you will know that you have reached the limit of in-breeding beyond which you should never try to go, but introduce another male bird with a certain amount of your own strain in him, when you will be able to go on again as successfully as before.

SCALE OF POINTS

Lacing	..	..	..	..	25
Comb, face and lobes	..	..	..	..	20
Ground colour	..	..	..	..	15
Tail ..	..	..	..	..	10
Type	..	..	..	..	10
Weight	..	..	..	..	10
Condition	..	..	..	..	10
					—
					100
					—

SERIOUS DEFECTS

Single comb; sickle feathers or pointed hackles on the cock; feathers on shanks; legs other than slate blue; other than four toes; any deformity.

CHAPTER X

SILKIES, NANKINS, JAPANESE, AND POLANDS SILKIES

OWING to its size, the Silkie has been included in this work. It is a question, however, whether it should have been, for it is not considered a Bantam, and when exhibited must be shown amongst the big breeds and not amongst the Bantams, unless of course there are classes specially for it. Many take it for a Bantam, and this accounts for so often seeing it shown in Bantam classes. By the remarks which follow it is evident that my late colleague, Mr. Proud, was of the opinion that the Silkie should be classed as a Bantam. In that opinion he was wrong.

Apart from its useful qualities as a sitter and mother when crossed with the Buff Pekin, the



Photo]

[Sport and General.

WHITE SILKIE.

Bred and exhibited by Mrs. Campbell.

Silkie hen is a decidedly pretty variety, with its silken plumage more resembling hair or down than feather, its skin, face and legs are black or mulberry; still there are the red-faced variety as well, but for my own part I prefer the mulberry-faced breed, as the contrast is so marked. The Silkie hen is also an excellent winter layer of tinted eggs, very docile, and one of the hardiest and easiest of all the Bantam tribe to rear. If you want a variety where accommodation in your backyard is at a premium, and one that will provide your children with a nice cream-coloured egg when the snow is on the ground, no other variety

turquoise-blue or mulberry ear-lobes, and legs and feet lead. It is a fine-toed breed. In breeding, only one pen is required, as it breeds very true to type and colour.

SCALE OF POINTS

Texture of plumage	..	..	30
Head	..	..	30
Type	..	..	20
Colour	..	..	10
Legs	..	..	10
			100



NANKIN BANTAMS.

The property of Mrs. Cross.

NANKINS

are a variety that appear to have almost dwindled out of existence.

It is very seldom now that we find a Nankin exhibited. Even at the big shows classes are not provided. From this it would seem that the day of the Nankin has gone, never to return. There is very little to add to these remarks, for the breed seems as scarce now as when the above notes were first written. It is very rare indeed that a Nankin is seen in the show-pen. This is perhaps a great pity, as it is one of the oldest of the Bantam tribe.

The Nankin cock is a very deep cinnamon or reddish-buff all over, except tail, which shows more or less black, the less the better, of course; legs and feet blue, comb, face and lobes red, comb single, eyes red. The hen is similar to the cock in all points except body colour, which instead of

being cinnamon should be a rich buff.

In breeding Nankins care should be used in selecting only those birds which are the soundest colour throughout, especially in flights and tail; any showing white should be rejected from the breeding-pen.

SCALE OF POINTS

Colour	..	..	..	25
Leg-colour	..	..	..	15
Shape	..	..	..	15
Size	..	..	..	15
Head	..	..	..	10
Style and carriage	..	..	..	10
Condition	..	..	..	10
				100

of Bantam will answer your purpose so well. The only drawback is her love to become broody.

On the other hand, if you desire to breed the very best of sitters and mothers to rear your Bantams, by crossing with the Pekin you will have the best natural incubator and rearer, one that can be relied upon at all times. They never run from their nest, but will sit till further orders, and very seldom do they crush a chick.

Silkie are apt to become scaly-legged, but in the majority of cases it is the result of uncleanness, and it is the same in nearly all the feathered-legged varieties.

The Silkie is seen in at least three colours, the white, black, and blue. It should have a black eye, mulberry face, small crest,

JAPANESE

have greatly increased in popularity during the past two or three years, thanks mainly to the exertions of a few of the chief breeders and exhibitors of this charming, and yet quaint, variety. At the chief shows no classes fill

is needed in breeding, as both exhibition cockerels and pullets can be bred from one mating. Japanese are amongst the oldest varieties of Bantams, but perhaps not so easy to rear as some of the other breeds. They are at their best from their second year onwards, being then better in shape, carriage



JAPANESE AND FRIZZLE BANTAMS.

The property of Major G. T. Williams.

better than those devoted to Japanese, and this healthy revival has been very pleasant to witness, and I trust it will be a lasting one, as it is a variety very suitable for the novice, no faking or trimming being necessary, and another great advantage is that only one pen

and length of feather than in their first season.

There are several colours, White, Black-pointed Whites, Brown-reds, Blacks, Greys, and Buffs being the principal ones, and of these the Black-pointed Whites, Whites and Blacks are the most popular.

The body is white and the tail black; the sickles of the cock, also side hangers, should be edged with white; face, comb single, and lobes red; legs and beak yellow, four toes, the wing primaries and secondaries black in inner web. Blacks are a sound black throughout and in all other respects are the same as the Whites. The Grey cock should be silvery white on back, wing-bow, and hackles on a black ground; the hen finely laced with silver, the lacing to be clear and distinct; comb, face, lobes and wattles same as above; eyes red, legs and beak yellow. Buffs should be what their name implies, a sound buff throughout, including tail and flights. Japanese, as a rule, are not at all difficult to breed and rear providing the ground is dry, and they breed true to colour and type.

SCALE OF POINTS

Colour of plumage	15
Head, and carriage of head and neck	15
Size, shape and carriage of tail ..	15
Shortness of legs	25
Colour of legs	5
Size	5
General shape	10
Condition	10
	100



Photo]

["Feathered World."

WHITE POLISH BANTAM HEN.

Exhibited by Mr. B. Wilkinson.



Photo]

["Feathered World."

BLUE JAPANESE BANTAM COCK.

The property of Major G. T. Williams.

SERIOUS DEFECTS.—Comb other than single, or in cocks falling over; shanks other than yellow, except in Blacks, Blues, Brown-reds or Birchen-greys; foul feathers in plumage, white in ear-lobes, long legs; long back, wry-tail, or low carriage of tail; any deformity.

POLANDS

are a variety that do not make many friends. Why this is so, I am at a loss to tell, unless it is the difficulty of breeding them true. They are a very pretty and taking variety. Unlike the larger breed of Polands, the most popular colours are the Whites and the White-crested Blacks, with here and there a Buff or a Gold and Silver; whereas in the large breed of Polands the favourite colours are the Silvers and Golds. Polands are doubtless a cross between the large Poland and the several breeds of variety Bantams, such as the Sebright and the Rosecomb, both Blacks and Whites.

If I were going in for breeding Polish I should use a big-crested Polish cock of the large variety, the larger the crest the better.

To produce Silvers I should of course use a Silver Poland cock and Silver Sebright hens. The following season I should pick out the best pullets from this produce and mate them back to the sire, choosing the pullets that were the best in crest formation. The following

season I should be able to reduce the size considerably by breeding from brothers and sisters ; selecting those which excelled in crest, I should then expect a reasonable percentage of Polish Bantams the following season, and would continue to inbreed again for another year or two until perfected. Golds could be produced likewise by using a Gold Poland cock and Gold Sebright hens, and Blacks from a white-crested Black Poland cock and Black Rosecomb hens. For producing Buffs I should use a Gold Poland cock and Nankin hens.

Of course, the first season you would have quite a number of unfertile eggs, especially so at the beginning of the season. This could to a certain extent be remedied by using the smallest male bird you could find, but possessing the needful large crest, which is absolutely necessary. Having done this, you must not expect it to be all lavender, but with perseverance and a reasonable amount of patience your labours will ultimately be crowned with success.

SCALE OF POINTS.

Colour	25
Size and shape of crest	25
Muffling	10
Comb	10
Shape and carriage	10
Size	15
Condition	5
	100

CHAPTER XI


CUCKOOS OR SCOTS GREYS, AND BARRED ROCK
BANTAMS
CUCKOOS

THE Cuckoos, or Scots Greys as they are called across the Border, are one of the most interesting of all varieties of Bantams. They are also entitled to rank among the oldest of the fixed breeds. It is nearly fifty years since they were first brought prominently before the public by Dr. Boulton, of Beverley, who, with his famous Field Spaniels and Bantams, was much interested in colour-breeding, and published a pamphlet on the subject. Although he undoubtedly originated the variety in England,

there seems every reason to think that similar birds had been produced in Scotland some years before that, and were known, as they are to-day, as Scots Greys. The late Mr. Henry Beldon was very successful in variety classes with a pair of Dr. Boulton's birds, and also, it is asserted, with *Scotch* bred ones. The Cuckoo hen is an excellent layer, and they are very hardy and easy to manage. I have often wondered how it is they have not become more popular. Reproducing, as they do, fairly well to type, it requires no great experience to keep a bird up to average form.

If, however, the amateur wishes to try conclusions at the Palace or Birmingham with the few enthusiastic veterans, who annually fight their battles there, he must be prepared to devote much thought and attention to them. One sees frequently a bird at the shows hit off by the reporters in a single sentence with a certain fault or other. In spite of all precautions, white feathers will come in wings ; white tips to otherwise perfect feathers, and the most promising cockerels have a provoking way, when least expected, and at the last stage, of developing white or black sickles or brown saddles.

It is a rare thing to see a cock's tail remain perfectly sound for the second year when one does stick to him for the breeding-pen. Opinions



LACED POLISH BANTAM HEN.

The property of Major G. T. Williams.

differ greatly as to the exact shade of ground colour. Many of the Scotch birds are too dark and indistinct in barring; on the other hand, many of the English birds are quite the reverse, being too light and washed out.

What I like to see is a clear steel grey ground colour, with close narrow distinct black barring. The greatest fault I have found with many birds successfully exhibited is that the barring is only black in the centre and not right across the feather. It is, in short, a modification of "mooning" rather than barring, and though the general effect to the eye produced by such a bird standing in the pen is much the same, when carefully examined in the hands it will be found to be absolutely wrong in principle.

The legs should be very fine in bone, quite free from feathers, and as white as possible. Spotted legs I do not greatly object to, but yellow or dark I cannot tolerate. The one great failing in Cuckoos is to keep the deaf ears red. Nature seems to object to it and continually endeavours to reassert itself. My advice to remedy this grave defect is to pay more attention to ears. It is interesting to note in passing that almost every black pullet produced from Cuckoos—and black pullets are produced in nearly every brood of good-coloured chicks—show white in deaf ears. You will invariably find that where you have one

or two black chicks in a brood, the remainder will be good-coloured ones, whilst if there are no blacks it is very probable the majority of the brood will be washed out in colour. So do not be disappointed when black chicks put in appearance providing they are not more numerous than the barred ones.

By far the worst faults of the Cuckoo Bantam of the present day are that they are yet too large and coarse in bone and lack the style and carriage we should like to see. In the endeavour to get them improved, we see birds exhibited decidedly gamey, short and narrow in feather, and too long in leg. On the other hand, we see birds much too long and narrow in back, and, worse still, with squirrel tails. The Cuckoo should be a short bird, broad across the back, full feathered, with the chest well expanded, the same as the Black Rosecomb, the head erect, and flowing tail well thrown back.

In mating up your breeding-pens of Cuckoos it is scarcely necessary to use two pens, as it has been found by our best breeders that the best cockerels are bred from the same birds as the best pullets of the same year.

The one thing needful in mating is never to fix a fault. Never, under any consideration, use stock birds that are white in ears or have feathers tipped with white. If your birds show a tendency to light ears, do your best to stamp out this grave fault by using a male bird that is absolutely sound in his ears, and which has been bred from sound-eared birds; and the following season use only the best of his produce, mating the sound pullets back to the sire; by doing this you will be remedying the defect, and building up your strain. Use those birds whose barring goes right across the feather (not just tipped), and as fine in barring as possible, and have the ground colour a sound steel grey, and the flights and tail as free from white as possible, you will not then get far wrong. Keep to one strain, and especially one which you know to be reliable, for in breeding for colour, as in Cuckoos, it is absolutely necessary to do so.

Keep a strict account of the pedigree of your stock, so that you can see at a glance how each bird is bred, and thereby recall to your mind's eye the chief characteristics of the bird, and whether such and such a fault is inherited or a mere accident, and mate him or her accordingly. There are plenty of thoroughly reliable stock birds which are not fit to grace a show-pen through having some glaring fault, but which are invaluable as breeders to those who are



SCOTS GREY BANTAM PULLET.

Exhibited by Mr. R. Fletcher Hearnshaw.

fully acquainted with their pedigree and how produced ; yet to anyone else these same birds would be useless, simply because they would not know how to mate them to remedy their existing faults. There is one thing about Cuckoos I feel certain of, and that is, upon no consideration in the effort to improve them should crossed blood be introduced. There is no breed which could possibly be used which would not damage them. On the other hand, as I have said in a previous chapter, they have been used freely in the production of new breeds of Variety Bantams. Strive to improve them by inbreeding ; fix for your standard the shape of the Rosecomb or Sebright, endeavour to keep them small and low on the leg, the leg to be as white as possible, carriage sprightly and erect, and steer clear of white in wing, tips of feathers, and tail. In point of colour, aim for the medium shade with a clear steel ground colour and rich, glossy black barring.

SCALE OF POINTS TO BE DEDUCTED

Bad head and comb	10
White in ear-lobes	10
White, black or brown feathers in wings and tail	10
Bad colour and barring	25
Bad feet and legs	10
Bad shape and carriage	15
Incorrect size and weight	10
Want of condition	10
	100

BARRED ROCK BANTAMS

Although seldom found in the show-pen, except where classes are provided for them, I do not think any other variety of Bantam offers a much better opening for a beginner than Barred Rock Bantams, and once they are firmly established they will undoubtedly find many friends, the same as the large breed of Barred Rocks, which to-day still holds its own in the soft-feathered varieties. It is only during the past eight or ten years that the Barred Rock Bantam has been introduced to the Fancy, but still they hardly make that headway one desires to see. The best of those which have come under my notice have been, for a new venture, decidedly good, although not so clear in ground colour and markings as we should have liked to have seen them, although the size and shape left nothing to be desired. There is no doubt whatever that by the introduction



SCOTS GREY BANTAM PULLET.

The property of Mr. J. McCrae.

of clearly marked Scots Greys the colour of the Barred Rock Bantam could be greatly improved, using those that are clear from dark spots on the legs, which we invariably find in Scots Greys. Still, by doing so, there is a great danger of losing the Rock type so essential in this variety. Therefore, great care should be exercised in introducing the Scots Grey.

Now I will proceed to give my idea of how to commence to breed Barred Rock Bantams from breeds other than Rocks.

Having obtained a couple of distinctly Barred Scots Greys, of the lighter shade, and which at the present time do not find favour with Scotch breeders, consequently would be cheaper and easier to obtain, and at the same time more suitable for the purpose—I should look round to find a cock the same colour, but with yellow legs, this being absolutely necessary, as the hens I am about to breed from are white-legged.

To meet these requirements I can find nothing better than the Cuckoo Pekin, which possesses the requisite barring and also yellow legs. Years ago it was quite possible to pick up yellow-legged Scots Greys, but these being an eyesore to Scotch breeders, were always killed off when very young, consequently the yellow leg was

entirely stamped out. I fancy I hear some of my readers remark, "Yes, but the Cuckoo Pekin has feathered legs." Quite so; and to remedy this grave defect as far as possible under the circumstances, I should try and get a Cuckoo Pekin as scantily feathered on leg as possible, at the same time I should require the leg colour to be a good rich yellow.

In making new breeds we cannot have everything cut-and-dried ready to our hand, but must use those which come nearest to our views of what is required. Naturally, the first season we should expect a fairly good percentage of feathered-legged ones, still not so heavily feathered as the sire, and in all probability there would be several perfectly free from feather on shanks, and these I should certainly retain as breeders for the following season. I should select the best cockerel, richest in leg colour, and mate him back to the Scots Grey hens, and if I could also mate up a small pen of, say, a cockerel and two pullets out of the first season's result, all the better, especially so if they were absolutely smooth in leg; but still I should not object to a few feathers on leg if the markings and type were there. You could then dispose of your Cuckoo Pekin, as you would have no further use for him. Having established the yellow leg, I should then proceed to improve the ground colour and markings. With this object in view I should select a cockerel free from brassiness on his back and one as free from white in tail as I could find. The clearer you have him in top colour the better he will be, either as a cockerel or pullet breeder. These two faults are very important ones when mating up your pens, and should always be borne in mind. By using this precaution you would save a lot of time and trouble, as it is an utter impossibility to breed clear-marked birds from a bronzy-backed male bird.

The hens to match the cock for pullet-breeding should be those clear and distinct in barring, with a nice steel-grey ground colour; the barring must be pure black, not a brown-black.

For cockerel breeding you would require the hens fine in barring and darker than those for pullet breeding; still, it is very important that the barring be a rich sound black and the ground colour not cloudy, otherwise your cockerels will come bronzy, which to-day is a very prevalent failing in Barred Rock cocks. Always strive to produce the clear steel ground colour, with rich black barring

combined with a bright orange-yellow leg and beak, then you will not get far wrong.

Do not expect 50 per cent. of flyers the first season, or yet the second; be content with three or four really good ones, stick to these, and inbreed, and with a little patience there is no reason but that in three years you will be able to produce Bantams equal in colour and type to the best Plymouth Rock of to-day.

SCALE OF POINTS

Colour and barring	..	..	25
Shape and carriage	..	..	15
Size	..	..	15
Head and comb	..	..	10
Ear lobes (red)	..	..	5
Legs and feet	..	..	10
Sound tail	..	..	15
Condition	..	..	5
			100

Faults to be deducted accordingly.

CHAPTER XII

FRIZZLES

WE have now come to the most quaint and charming of all Variety Bantams, viz. Frizzles, which, notwithstanding the fact that they have been exhibited in England nearly forty years, have failed to "take on" in the Fancy. This is probably on account of the difficulties experienced in rearing the chicks; as the variety has been greatly inbred in order to obtain the type and character so essential in Frizzles, the stamina of the breed has been greatly weakened.

There is great diversity of opinion as to what country this breed originated in. Some say Japan, others Friesland, whilst some ideas of their origin are both too numerous and too funny to enumerate in these notes. However, their nationality matters little; they are here, and having been here so many years are likely to remain, especially seeing that both judges and fanciers are gradually beginning to appreciate their charms.

Although they have invariably to compete against many varieties, even at such shows as Dairy, Palace, and Birmingham, they generally give a good account of themselves, as reference to reports will show.

The most trying time with Frizzle chickens is the period of getting their first feathers. At this time they are difficult to rear, but when the ground is dry and sandy successful results can be obtained. Having got their first feathers they will be found quite as hardy as most varieties.

Many persons imagine that curling-irons play an important part in preparing Frizzles for exhibition, and in some cases such is the case. I remember once seeing a lady manipulate with the curling-irons on a white Frizzle hen for two hours one Sunday night, and although I have often heard it said that "better the day, better the deed," it was not so in this case, for this exhibit just managed to struggle into



BLUE FRIZZLE BANTAM HEN.

Bred by Major G. T. Williams.

fourth place. Then, again, I remember some years ago, either during the Palace or Dairy, the latter I believe, a certain exhibitor had occasion to attend the hairdresser's shop near the show, and on entering was surprised to see a brother fancier, one of the lilywhite innocent ones, too, holding a wee Frizzle pullet on the back of a chair whilst Mr. Hairdresser was assisting nature to curl the feathers. On this occasion their united labours were rewarded with success.

There are few, very few, if any of the varieties of poultry that cannot be materially improved by artificial means, and Frizzles may probably be no exception to the rule. Still, in many breeds of to-day science in breeding has played such an important part as to leave no necessity to improve. So it is with this quaint variety; when carefully bred on scientific lines with a view to perfect the type and character, and

given a thorough good wash and a nice bright fire to bring out the desired curl, there is no necessity for curling-irons to "fake"—I beg pardon, I mean, to assist nature.

Frizzles appear at their best on either a warm summer day or a sharp, clear, frosty day, but the slightest fog or rain deprives them of their beautiful curls at once, the same as it does the ladies' curls when not in papers. This is one great drawback (to the Frizzles, I mean) when they have to be sent long journeys in the damp winter months.

Frizzle Bantams throw a large percentage of plain-feathered chickens, utterly useless for exhibition, but are of invaluable service in the breeding-pen; in fact, really good specimens are rarely produced without them.

The curled hens are excellent mothers and sitters, but the plain-feathered ones are not to be depended on in either capacity. This seems strange, but nevertheless it is quite true. There are now many colours of Frizzle Bantams, including several colours as seen in the O.E. Game, such as Spangles, Blue-reds, and blue-tailed Wheatens. Still, Whites, Blacks, Buffs and Blues remain the popular colours. The plumage should be broad, long, and crisp, each feather curling towards the head, with frizzling close and abundant. The colour of the plumage should be uniform in Blues, Blacks, Buffs, and Whites, and other colours as that of O.E. Game. Leg and beak colour should be yellow in Whites and Buffs, black or blue in Blues and Blacks, and in other colours as that of O.E. Game. To breed Frizzle Bantams it is advisable to have two breeding-pens; this is only necessary on account of getting *curl*, as one breeding-pen if properly mated will produce both good cockerels and pullets.

No. 1 pen should consist of a hard-feathered, well-curved cock, short in back, full in frill and breast, with short legs of the standard colour; mate to him two small well-curved hens and two plain feathered hens or pullets, whose appearance should be as round and as squat as possible.

No. 2 should consist of a cock similar to above, but plain in feather and very full feathered; to him mate three or four well-curved hens, the closer in curl the better. The plain pullets from No. 1 pen I should use with the best curled cock the following season.

It is no use trying to rear Frizzle chickens in February or March, it will only be labour in vain. The months of May and June are most

suitable for this variety in England, especially in the northern counties, and not earlier than April in the south.

SCALE OF POINTS

Curl	..	..	..	..	30
Colour	..	..	..	..	25
Weight	..	..	..	..	10
Condition	..	..	..	..	10
Type	..	..	..	..	25
					100

SERIOUS DEFECTS.—Want of curl; narrow feather; white lobes; long tails; drooping combs. Disqualifications.—Other than single combs; other than four toes on each foot; feathered legs.

CHAPTER XIII

BUFF ORPINGTONS, ROCKS AND LEGHORNS

HAVING described fully the Barred Rock Bantam, I will now go on to show how it is possible to breed the fashionable Buffs.

The nearest approach to Buff Orpington Bantams came under my notice on a visit to the yards of Mr. John Wharton during May, 1898; the parents of these I learned had been imported across the pond, along with some Buff Wyandotte fowls from the United States. These were very small, short on leg, and a very level buff for a new breed, and although free from feather legs they possessed quite a bluish tinge in leg colour. This, I informed Mr. Wharton, was a very serious fault, and that if he could only obtain the necessary white leg he would undoubtedly "strike oil," as the Yankees say.

The next time I saw Buff Orpington Bantams (if you will excuse the name) was at Carlisle Show, November, 1899, where a class had been provided for them, but instead of Buff Orpington Bantams they reminded me of "the Boer Army," a mixture of all nationalities, but never a trace of Buff Orpington could I find, and it must have puzzled the judge not a little to come to a decision in his awards.

The only rational way that appeals to me for the production of Buff Orpington Bantams, is to get a very late hatched Buff cockerel of the large breed, as small as possible and sound in colour, and mate him to a pale-legged Buff

Pekin, as scantily feathered as possible. The following year I should inbreed from the produce of this pair, and should there be one or two pullets extra good in leg colour and free from feathering that I could spare (after mating up a small pen of a cockerel and two or three of his sisters), I would put the former pullets back to the father; afterwards I should inbreed and breed late in the season for some three years, by which time I could reasonably expect a fair percentage of exhibition specimens.

The necessary requirements would be the same as in the larger breed of Buff Orpingtons, viz. pure rich orange or golden buff; clear buff in flights and tail, no white or black feathers. The former I should consider a fatal defect in the show-pen. Legs, perfectly white and free from feather; shape, same as in Orpingtons, viz. low on leg, deep square body, and short in back; comb single or rose; lobes perfectly red, any white in lobe to count heavily against the bird in the show-pen; size as small as possible consistent with shape.

SCALE OF POINTS

Colour (sound level buff, including flights and tail)	..	..	25
Shape and carriage	..	..	20
Pure white smooth legs	..	..	15
Sound red ear lobes	..	..	15
Size	..	..	15
Condition	..	..	10
			100

Faults to be deducted accordingly.

DISQUALIFICATIONS.—Feathered legs, white in tail and flights, pulled tail or shanks, white in lobe, leg colour other than white.

Weight.—Cocks not to exceed 24 oz., pullets 18 oz.

ROCKS (OTHER THAN BARRED) AND LEGHORNS

Buff Rocks used to come next to the Barred variety in popularity, but this colour never took on; and it has declined to such an extent that now it is very rarely seen. This is rather a pity, for it was a very taking variety, and for type it equalled any of the Rock tribe. In the early days of its existence, Mr. Jones, of Middlewich, was an enthusiastic breeder, and Mr. Hipkins, of Tipton, who I believe took them up later, was a very great force, but owing to business engagements he had to give them

up. In recent years Whites have been fairly numerous, and some very nice specimens have been exhibited. They come next to the Barred in public esteem, though they are hardly ever likely to oust that colour from premier position. The White Wyandotte has, in some strains, played its part in the production of this colour. Blacks are occasionally seen, but this colour is not, I think, very extensively bred. Soundness and uniformity of colour, along with type and carriage, are important factors in this breed.

Leghorns are a breed that has never really caught on, and at the present time I should say there are few birds in the country. At times it has been seen in many colours, comprising Whites, Blacks, Reds, Blues and Buffs. Both the Rock and the Leghorn present a capital opening for a beginner, for besides their possibilities in the show-pen, they are above the average in utility qualities.

SCALE OF POINTS

Colour, sound throughout, including tail and flights	25
Shape and carriage	20
Size	15
Sound ear-lobes	15
Rich smooth legs	15
Condition	10

100

Faults to be deducted accordingly.

DISQUALIFICATIONS.—White in tail and flights, pulled tails, leg colour other than yellow, and in Buff Rocks white in lobe.

Weight.—Cocks not to exceed 24 oz., pullets 18 oz.

CHAPTER XIV

MINORCAS, ANDALUSIANS AND SPANISH


SINCE the last edition of this book appeared, Minorca Bantams have made rapid headway, and, with the formation of a Club to further their interests, they should make still more progress. Some excellent specimens, both as regards head points and size, are now staged. A little more length of back would be an improvement, the shortness in this respect being no doubt due to the Black Rosecomb cross. The Club caters for both colours, the Black and the White, but the latter colour is not very often seen. The breed appears to

have a bright future, and any one is hardly likely to regret taking it up.

VALUE OF DEFECTS IN JUDGING MINORCA BANTAMS

Defects in face	15
Bad shape or twisted comb	15
Over size	15
Wrinkled, folded or stained lobe	10
Defects in colour	10
Want of condition	10
Want of style and symmetry	10
Too light legs, eyes or beak	8
Crooked breast-bone	7

100



MINORCA BANTAM COCK.

The property of Mr. T. Bowker.

SERIOUS DEFECTS.—White in face; wry or squirrel tailed; feathers on legs; other than single combed; coloured plumage other than black or white in the two colours; other than four toes; legs other colour than black or slate in the Black variety, or white in the White variety.

To produce Andalusian Bantams I should use a small, distinctly-laced Andalusian cock, not too dark, a shade light would be all the better, but the *lacing* must be *perfect*—this is very important. I should then inbreed from the single-combed produce of this hen until I

obtained the desired type and colour, which should be a miniature of the Blue Andalusian. Of course, it is quite possible to get good Minorca Bantams from this cross as well. Still, I should prefer to breed Andalusians only, as they are much prettier and command a better sale, for the Minorca Bantam looks too much like a single-combed Black Rosecomb.

The Andalusian Bantam is probably now extinct, at least it is many years since any were exhibited. It is a pity it did not make more headway when it was first introduced, for there is no denying its beauty.

The Andalusian Bantam should represent the large Andalusian in colour, shape, and carriage. It should be compact and erect, rather square in build, fair length of back, tail full and flowing, and carried well back; comb, single and well serrated, erect in cocks, and folded in hens; face, smooth and red, any white in face being a serious fault; lobe, large, smooth, and white, almond shaped, the larger the better, consistent with a sound red face, which are difficult points to obtain together, for invariably when a bird excels in size of lobe he is liable to run white in face; legs, not too short, and dark slate in colour.

The colour of the cock should be: Hackles, back, and wing-bow a soft shade of bluish black, the remainder of the plumage being medium slate colour or inclined to grey, each feather distinctly laced with black. The hen to be medium blue-grey in ground colour, one even shade all through, and each feather distinctly and evenly laced with black. The neck hackle should be a bluish black.

SCALE OF POINTS

Colour and lacing	..	..	35
Comb and lobe	..	..	15
Size and weight	..	..	20
Shape	..	..	10
Condition	..	..	20

100

DISQUALIFICATIONS.—Wry back or tail, folded comb in cocks, ear lobes other than white.

I should like to see the Spanish Bantam more popular, and remember judging at Kendal Show some years ago, where Spanish Bantams, little beauties too, real miniatures of the present-day Spanish, with smooth white open face and glossy black plumage, stood first and second. How they were produced I know not,

but I should think by using a small Spanish cock, excelling in face, and mating him to a big-lobed Black Rosecomb hen, which, as a rule, you will find are more or less white in face, the more the better in this case, for your requirements. If she is a bit large never mind, as you would then get more fertile eggs than from a small one, therefore more chicks to choose from for your next year's mating. The pullets that were best in face and possessing the necessary single comb I should put back to the sire and breed as late as September the following season, and afterwards inbreed for the next two or three years, brothers and sisters, etc., always having the head points and large white open face as your guide in mating up, rather than size, as this can always be reduced by inbreeding and breeding late in the season. The Spanish Bantam cock should have single comb carried erect, large open face, the wattle to be as white and smooth as possible, and without any break in the face, eyes dark, plumage raven black, not too low on leg, medium length of back, tail full, flowing, and carried fairly well back, legs as near black as possible, the hen to be the same, with the exception of comb, which in all the three varieties should be folded.

SCALE OF POINTS

Face and lobes	..	..	..	30
Size and weight	..	..	..	20
Comb	..	..	..	15
Colour	..	..	..	10
Condition	..	..	..	15
Shape	..	..	..	10

100

DISQUALIFICATIONS.—Wry tail, or wry-backed, any coloured feathers except black.

CHAPTER XV

HAMBURGH BANTAMS

WISHING to include a description of these Bantams, I bethought myself that I could not do better (seeing I had never bred these beautiful varieties) than write to Mr. John William Farnsworth, whom I knew had made a speciality of Silver Spangled Hamburg Bantams for many years, not only breeding them, but also exhibiting them successfully at a few shows where the classification was at all suitable.



OLD ENGLISH GAME BANTAMS.

1. SPANGLE COCK.

2. SPANGLE HEN.

3. DUCKWING COCK.

4. DUCKWING HEN.

5. BLACK-RED COCK.

6. WHEATEN HEN.


7. BLUE-RED HEN.

8. BLUE-RED COCK.

To my request for information for the readers of *The Feathered World* as to how he started and eventually perfected the Silver Spangled Hamburg Bantam, Mr. Farnsworth wrote me a thorough Fancier-like reply, which I cannot do better than give in his own words:

"The following I do not say is the right way to go about the work, but it is the line pursued in producing my birds.

"My accommodation has always been limited, hence anything but Bantams in the fowl line were out of the question. Born with the hen-fever of a permanent character, a general liking for anything small, and a great admirer of the 'black and white' combination, the Silver



Photo]

["*Feathered World*."

SILVER HAMBURGH BANTAM.

Exhibited by Mr. G. Gresswell.

Spangled Hamburg Bantam, although only existing in my mind's eye, stamped itself at once to be *the* very thing. A really good but undersized Silver Spangled Hamburg cockerel of notable descent, having been mated to a couple of rather large Silver Sebright hens, resulted in my being, the following year, in possession of a fairly good Mooney cross cockerel, which was mated to a somewhat large blue-legged White Rosecomb hen, a Black Rosecomb hen, and his sister, a Mooney cross pullet. Another similar pullet was bred to a pure White Rosecomb cock.

"The remaining Mooney cross pullet, the smallest of the three, was put back to her sire, the Hamburg cock, the services of the Sebright hens being at once and *for ever* dispensed with.

"These three pens produced a number of odd-looking chickens, of various markings, but a very good selection was obtainable therefrom, some of which were bred together, others were mated with White Rosecombs and Black Rosecombs, to improve size and spangling. By continued perseverance I produced some grand little birds in every respect but tail, which I could not get satisfactory. Eventually I bought a Silver Spangled Hamburg cock with a *perfect* tail and but $3\frac{1}{2}$ lbs. in weight (of the same strain as the original Hamburg cock), which, mated to a couple of my largest Spangled hens, gave me some chickens with the desired tail, but, of course, oversized.

"These, bred together, with care in selection, for a few seasons, enabled me to pen three birds at Basingstoke show, and I was awarded first prize in the A.O.V. cock class, and second and third prizes in the corresponding hen class—a, to me, very gratifying result. Since then I have but once missed the money when exhibiting them, and that at the Midland Counties Bantam Show, at West Bromwich, where in seventeen entries I got h.c.

"I have, of course, had the usual amount of *trouble, expense, and disappointments* in bringing them into form, but I consider them without doubt one of the handsomest Variety Bantams out at present, and *The Feathered World* and other Fanciers' papers have all spoken most favourably of them."

Such is Mr. Farnsworth's method of producing Silver Spangled Hamburg Bantams, which are without doubt one of the prettiest of all Variety Bantams, and the production of which presents no difficulties whatever. With ordinary luck and careful selection each year, by inbreeding after having obtained the desired colour, and by breeding as late as June to reduce the size, there is no reason why exhibition specimens should not be produced in about four years. The only difficulty I can see would be in keeping the length and flow of tail feather, which, to a certain extent, would be lost in breeding very late.

Colour and length of feather should be your guide in mating up your breeding-pens, rather than size, as size can always be remedied, and very materially so, by feeding and inbreeding.

Gold Spangles could easily be bred in the same way by using a Gold Sebright hen, but they are not nearly as beautiful as the Silvers, and I do not think for one moment that these would ever become popular, or pay for the time and labour.

GOLD AND SILVER PENCILLED

These varieties have now been added to the list of Bantam breeds, and when they have been got to anything like a state of perfection they cannot help but make many friends, as their colour and fineness of pencilling are really beautiful; in fact, they are likely to run the Spangles a close race for supremacy. I do not think it would take a great length of time if breeders of these varieties were fully determined to produce typical Bantams in every respect equal to their larger brethren.

The best plan to commence operations would be to mate up two or three small finely Gold Pencilled Hamburg pullets, rich in ground colour and good in head points, and mate them to a Gold Sebright cock, as fine in the lacing as possible, using a cock or cockerel possessing sickles, as some cocks have little or no sickles, and from these I think you would be able to obtain some very passable Gold Pencilled

Hamburg Bantams; and, by mating up the best cockerel back to the hens again the following season, and hatching the chicks between June and August, great improvement could be made the following year again by inbreeding with cockerels and pullets of this produce, selecting those with the richest ground colour, and finest pencilling; and the following year it would be advisable to inbreed again, and as late in the season as possible, but do not attempt to hatch the chicks before the month of May.

In breeding Silver Pencilled I should adopt the same method, using a Silver Sebright cock, one extra fine in lacing but good in head points, and mate him to the smallest Silver Pencilled Hamburg pullets you could lay your hands on; then the following season inbreed as before, mating the smallest and best-coloured cockerel back to the hens and one or two of the smallest pullets that show the correct pencilling, and the following year inbreed from brother and sister with the smallest and best-marked birds.

By paying strict attention to size and markings, and not hatching the chicks before the month of May, and between that month and July, there is no reason why typical Bantams of these varieties should not be produced in three years from the start; and, once they are produced, I do not think there is any other variety of Bantam that would find more favour on account of their exquisite markings and colour—especially in the pullets, which would be very handsome indeed.

Of course, there is one great point in the breeding of Hamburg Bantams which must not be lost sight of, and that is length of feather, for without this the cocks would be shorn of much of their beauty, although in pullets it does not matter so much; but there is nothing that sets off a Hamburg cock half so much as a long flowing tail with plenty of side hangers or streamers, as they have been called; and to acquire this it will be necessary to always breed from the best-feathered cocks and the longest-tailed hens which have good flowing neck hackles.

In breeding Hamburg Bantams, and especially when you have to use a certain amount of Sebright blood to reduce the size, you will be very liable to produce cockerels with hen-feathered tails—i.e. tails the same as the hen, with very little or no sickles; and,



Photo]

["Feathered World."

GOLDEN PENCILLED HAMBURGH BANTAM.

Exhibited by the Rev. M. L. Thorp.



PARTRIDGE WYANDOTTE BANTAM PULLET.

The property of Mr. Scott Miller.

although these as a rule will be the best-coloured birds and are all right as pullet breeders, they should never be used with a view of producing good cockerels. Therefore it will be advisable in breeding these two varieties of Hamburg Bantams to mate up separate pens for cockerel and pullet breeding. For cockerel breeding, let the male bird excel in length of feather, long sweeping sickles, and plenty of side hangers, and mate him to hens that have long neck hackles and plenty of tail, even though they be somewhat faulty in colour, although it will be necessary to have the cock as pure in colour as you can get him. In pullet breeding this length of feather in the cock is not desirable or necessary, but the hens or pullets must possess fineness of pencilling and rich ground colour in the Golds, and fine marking in hen in Silvers, with good white ground colour, free from any trace of yellow or creaminess, which will be the one prevailing weakness in this variety.

When making a start with the Hamburgs it will be best to aim at one variety first, and procure the pullets from some breeder who has made pullet breeding a speciality for years, and the same way with the cocks, then you are all the more likely to produce typical specimens in due time.

Although the standard is given herewith for Hamburg Bantams, I do not think it would be out of place for me to draw up one for these two varieties, which will be a help in enabling young fanciers to know the principal

points that are required in Gold and Silver pencilled Bantams.

The chief points are undoubtedly fineness and evenness of pencilling, the pencilling to run straight across the feather in a clean-cut fine bar, as narrow as possible, and this, together with a sound ground colour, should count 25 points. Then comb and head points 10, length of feather 15, type 20, size 15, and condition 15—a perfect bird to count 100. In reference to the size, for the first two or three years I think 18 oz. to 24 oz. should be allowed for cocks, and 15 oz. to 20 oz. for hens; but, of course, the smaller the better, colour and markings being equal.

SCALE OF POINTS

Head and comb	25
Lobes	10
Colour and markings (except Blacks)	15
Shape and carriage	10
Size and weight	20
Condition	20
	100

DISQUALIFICATIONS.—Wry tail, single comb, and deformity of any kind.

CHAPTER XVI

WYANDOTTES

SINCE this breed was first introduced some twenty years or so ago, no new breed has made such rapid headway in the Fancy as the Wyandotte, not only in perfection but also in public esteem. It can now hold its own against most of the old breeds, and it is one of the most popular breeds in the soft-feathered section. When the first edition of this book was published, the Partridge and the Silver-pencilled were the leading varieties of the Wyandotte family, and such excellent progress had been made that in colour and type they, at that time, were almost equal to the best specimens seen in the large breeds.

When I revised the last edition of this work, the Blacks and Whites were of comparative recent introduction, and I gave it as my opinion that the Whites were likely to rank as one of, if not the best of all the varieties of the Wyandotte. That opinion has proved

wonderfully true, for at the present time the White Wyandotte stands out as the most popular variety in its section.

In no other breed, with the exception of Old English Game and Belgians, is there the same choice of colour as in the Wyandotte, for besides the colours named, there are Blues, Buffs, Columbians, Spangles, Silver-laced, Gold-laced, Buff-laced, Blue-laced, etc. Of course, a good few of these colours are not very popular, and some of them are only rarely seen. The most popular colours are Whites, Blacks, Columbians, and Partridges. Partridges continue to hold their own well in a steady fashion, but Blacks, a variety that some years ago were about the most popular of the lot, now appear to be hanging fire, but this is probably only one of the minor effects of the war, and no doubt in the near future all the old enthusiasm will again be seen. Columbians have made splendid headway, and in the near future this colour is likely to go still further ahead. Blues and Buffs should make progress.

PARTRIDGE

The male bird should be identical with the large Partridge Wyandotte. The comb should be of fair size, fitting nice and close to the head, with the leader, or spike, as it is sometimes called, following the curve of the neck. The more work in the comb the better. The comb, face, wattles, and ear-lobes should be a bright healthy red, white in lobe being a serious fault, and this should be strictly guarded against when mating up the breeding-pens. We now come to the neck hackle, which should be a rich golden-orange, each feather having a green-black stripe down the centre, although many will be found with a brownish-black stripe, which is a bad fault. The saddle hackle should correspond as nearly as possible with the neck hackle, and the brighter in colour the better. The back from where the neck hackle falls to should be a rich, bright, dark red, and this should continue to where the saddle hackle commences. The wing-bar, breast, right through to the tail, should be a rich, glossy green-black; the tail should be the same colour, also the flight feathers of the wings. The wing-end, when closed, should be a rich chestnut or bay, legs and feet a rich orange-yellow, and the shanks free from any sign of feathering.

The most serious fault to be found in the cocks is white in flights and root of tail, and

this can only be detected by handling the bird very carefully and opening out the wings and tail. In shape the bird must be very cobby in build, full in chest, and not too long in back or too high on leg.

The pullet to match the exhibition cockerel I have described must resemble her lord and master in type, the cobbler in build and the smaller the better. The neck hackle should be striped with black, as in the cocks, but this is a most difficult point to get in conjunction with a proper shade of ground colour and the clearly defined pencilling in the feather. The ground colour of the pullet should be one even shade of light brown, and each feather should show distinct pencilling of green-black, with an outside edging of the same shade as the ground colour; the more defined and clearly outlined the pencilling the better all over, although it is somewhat difficult to get this clearness of pencilling on cushion, where quite 50 per cent. fail. The legs should be yellow, and the comb, face, lobes, and wattles a bright red, as in the cock. The pullet I have described is an exhibition specimen, and such should be used for pullet breeding, with a cock or cockerel of a shade or two darker top colour than the



Photo]

[*"Feathered World."*

PARTRIDGE WYANDOTTE BANTAM HEN.

Bred and exhibited by Mr. J. F. Entwistle.

exhibition cock, and one showing lacing on breast and fluff behind the thighs the better.

The exhibition cockerel, as I said before, should be as bright in colour as possible, and perfectly sound in his black on breast and fluff, and the pullets to mate to him for producing the best cockerels should be a very bright orange in neck hackle, but the striping must be as broad and defined as possible, and a rich, sound black, although they will be found very indistinct in pencilling—in fact, scarcely pencilled at all—and much darker in ground colour. Such pullets, although worthless as exhibition specimens, are valuable as cockerel breeders.

In breeding Partridges, it is almost a necessity to have two pens to give the best results, and if care is used in the selection of the stock birds as I have described a fairly good percentage of exhibition birds should be bred each season. Do not sacrifice colour for size, but try and maintain the nice, cobby, squarely-built type, purity of colour and pencilling; then you can easily reduce the size later by in-breeding and by hatching the chicks late in the season.

SILVER-PENCILLED


These followed the Partridge, and are identically the same in type, head points, and pencilling, but where the cock is orange and red in the Partridge, the Silver-pencilled is a silvery white, and the purer the white the better. The wing-bars, breast, underparts, fluff, tail and flights, black, as in the Partridge, legs and feet yellow.

The hen should be pencilled the same as the Partridge on a ground of steel grey, and the mating recommended for Partridge should be maintained in the Silver-pencilled in every detail.

This brings us to the

BLACKS

These are amongst the latest addition to the Bantam Fancy, being brought out in the year 1907, and were undoubtedly produced by a cross between the Partridge and the Black Rock Bantams, the latter being the sports from the Barred Rock Bantam, and this is one of the best ways of producing them. The Black Wyandotte Bantam should be a rich green-black from head to tail, including the flights and ends of the wings, also tail and sickles, the latter being the most difficult point to get



BLACK WYANDOTTE BANTAM HEN.

The property of Mr. H. Phillips.

sound, as the majority of Blacks fail at the root of the sickle. In shape they should be cobby, squarely built, and not too high on the leg. The comb should be the same as described on the Partridge, and the face, wattles, and ear-lobes should be a bright, healthy red, white in lobe being a serious fault. The beak and legs should be yellow, although it is a very difficult matter, and will be for some time to come, to get a good yellow beak. The under-colour in exhibition birds should be black to the skin, although many would be found to be quite white; and although this is a serious drawback in the show-pen, such birds, as a rule, are useful in the pullet-breeding pen, but for breeding cockerels you should have the male bird a sound black to the skin, including the neck hackle, where quite 50 per cent. are inclined to go very light at the root of the feathers. The surface feathers should be a rich green glossy black, free from any trace of bronze or rainbow shade; in fact, any colour other than black should count heavily against a bird in the show-pen.

The most serious faults should be legs other than yellow, white in flights, tail or lobe; and the minor faults, white in under-colour, faulty shape, too leggy or spotted legs, although of a yellow ground colour. The breeding of Blacks should not be so difficult as the other two varieties I have just mentioned, and it would be quite possible to breed exhibition



BLACK WYANDOTTE BANTAM COCK.

The property of Mr. H. Phillips.

specimens of both sexes from the same pen by judicious mating. In breeding cockerels it is absolutely necessary to have the male bird a pure black to the skin, also sound in flights and tail, but should you have a bird that shows plenty of green in the black and is sound in under-colour, but a bit red on the saddle and neck hackle, you must not discard this bird, as he will be found to be a valuable pullet breeder, but must not be used for cockerel breeding. The pullets to mate to a sound-coloured Black cockerel should be as rich a green-black as possible, and should also be good in head points, and excel in type, but it is from the female side that the shape of the offspring is influenced. For pullet breeding you can use a cock bird showing a little red on the saddle and hackle, as I have described, or one that is a little light in under-colour, but see that the hens are sound and good in head points, also rich in leg-colour. To obtain the best results from one pen it will be best to use a sound-coloured male bird, rich in green in top-colour, sound flights and tail, with rich orange-yellow legs, and if he be a little light in his under-colour, it will be no great objection if he excels in other points. He should be

mated to good-shaped hens as green in colour as possible, and possessing good legs. When using the Black Rock Bantam hens to a Partridge cock, the latter should be as dark as possible in colour, and from this mating you should be able to get two or three good birds the first season, although no doubt, many will come with single combs, and though useless in the show pen as Black Wyandottes, will be good material for next year's breeding put back to a Black cock, providing they are yellow in leg or show a fair amount of yellow tinge in leg-colour.

WHITES

This colour is one of comparative recent introduction, having been launched on the Fancy about fifteen years or so ago. It is a charming variety, and its popularity is not surprising. It looks like retaining for a long time the position it at present holds. Splendid progress has been made in perfecting it, and many almost ideal specimens have been penned during the last few years. The fact that there is little or no necessity for double mating is a great point in its favour, and for that reason, it can be highly recommended to the amateur's notice. Type, size, carriage and purity of colour, with neat, yet stout heads and good yellow legs are the chief features, and in mating up the breeding stock, those are the points to bear in mind.

COLUMBIANS

This is a most pretty and useful variety, and one that is making steady progress. Probably during the past two or three years it has come on as well as any of this breed. Very few really good cocks or cockerels have been seen, but some charming hens and pullets have been penned. Many exhibits have been lacking in type, and to my mind that is one of the most important features in any breed. No doubt in time that defect will be overcome, but it behoves breeders not to lose sight of it when mating up the breeding-pens. In colour of plumage the Columbian Wyandotte practically resembles that of the Light Brahma, and in breeding most of the points laid down for the latter apply equally to the Columbian. Type, soundness of hackle striping, and purity of top colour are the great features to bear in mind.



WHITE WYANDOTTE BANTAM COCK.

The property of Mr. E. Whitaker.

OTHER COLOURS

Golds and Silvers, and Buff-laced and Blue-laced are very rarely seen, and probably the breeders of all four could be counted on the fingers of one's hands. There can be no denying the pretty combination of colours, but somehow they have failed to take on, and it hardly looks likely that they ever will. They seem to have followed the lead of the large varieties of those colours. Spangles are more often seen, and in time, these may become fairly popular. These are sound black in ground colour, with the tip of each feather distinctly tipped with white. They first originated as sports from the Blacks. A few useful Buffs have been seen very nice in colour, but as yet rather lacking in type. Buff Columbians, marked like the Columbians but with buff ground colour, are about the latest acquisition to the breed, but it is too early to say much about them. Messrs. C. & S. Ford exhibited one at the 1923 Dairy Show. Blues have been more or less bred for some years, but somehow, they have not struck the popular fancy. Probably they have been exhibited more during the past two seasons than ever before, and some very typical ones in shape have been seen. There is no doubt about it being a taking colour, but it is a difficult one to breed true to colour. The pullets are easier to breed true in this respect than the cockerel, as it is in the case of the big variety of this colour. A useful Cuckoo or two are now and again to be

seen, but this is another colour that fails to take on.

STANDARD OF PERFECTION OF WYANDOTTE BANTAMS

Cocks, 28 oz. ; Hens, 24 oz. ; Cockerels, 24 oz. ; Pullets, 20 oz.

GENERAL CHARACTERISTICS

COCK :—*Head and Neck*.—*Head*, short and broad. *Comb*, rose, firm and even on head ; full of fine work ; low and square at front, tapering towards the spike, which should follow the curve of the neck. *Face*, smooth and fine in texture. *Ear-lobes and Wattles*, medium length ; fine in texture. *Neck*, medium length, well arched, with full hackle.

Body.—*Breast*, full and round, keel bone straight. *Back*, broad and short. *Saddle*, full and broad, rising with concave sweep to tail.


Wings.—Medium size, nicely folded to the side.

Tail.—Well developed, spread at base, the true tail feathers carried rather upright, sickles of medium length.

Legs and Feet.—*Thighs*, of medium length, well covered with soft and webless feathers. *Fluff*, full and abundant. *Shanks*, medium length, strong but fine in bone. *Toes*, four in number, straight and well spread.

General Shape and Carriage.—Graceful and well balanced, resembling a Brahma.

HEN :—*Head and Neck*.—*Head, Comb, Face, Ear-lobes and Wattles*, as in the cock, but the appendages smaller. *Neck*, medium length, with short, full hackle.



WHITE WYANDOTTE BANTAM HEN.

The property of Mr. E. Whitaker.

Body.—To correspond with that of the cock.
Back, short and wide at shoulders.

Tail.—Well spread at base.

Legs and Feet.—As in the cock.

General Shape and Carriage.—To correspond with the cock.

SERIOUS DEFECTS FOR WHICH BIRDS SHOULD BE PASSED

Any feathers on shanks or toes. Permanent white or yellow in ear-lobes covering more than one-third of their surface. Combs other than rose, or falling over on one side, or so large as to obstruct the sight. Wry tails. Deformed beaks. Crooked backs. Shanks other than yellow in colour (except adult cocks


generally found in the brightest-coloured birds, judges must use their discretion in penalising according to the colour of the bird and the degree of the defect.

In the Female.—Light shaft to feathers. Red or yellow ground colour. Light under-colour. Smutty legs.

Standards of the various colours can be obtained from the Club Secretary, whose address can be had from *The Feathered World* Office if a stamped addressed envelope is sent.

BELGIANS.

It was, I believe, at the Crystal Palace Shows in 1911 and 1912 that our Belgian friends brought over an interesting collection of their



A QUARTET OF COLUMBIAN WYANDOTTE BANTAMS.

(Utility with Beauty.)

Owned by Mr. J. E. Cross.


and hens, which may shade to light straw colour). Feathers other than white in Whites. White in tail or any conspicuous spotting or peppering on ground of the feathers in Silver and Gold.

OTHER DEFECTS IN PARTRIDGE WYANDOTTES FOR WHICH BIRDS SHOULD BE PENALISED

In both sexes.—Crooked toes, the extent of the defect to determine whether a bird should be passed.

In the Male.—White in tail and at root of tail. Grey fluff. White flights. As these defects, together with light under-colour, are

favourite Bantams, and these birds naturally attracted much attention from fanciers and other visitors to those shows. Many charming varieties were exhibited, and certainly quite a good number of fanciers took them up, with the result that they were becoming quite popular when the Great War broke out. During the fateful years which followed Belgian Bantams declined rapidly, with the result that they were hardly ever seen except at a few of the larger fixtures, and at these the entry was few in numbers. Still, a few keen enthusiasts stuck to the breed, and during the past three or four years a revival has been quite apparent. It is to be hoped that now things look brighter



A PRIZE-WINNING BELGIAN BANTAM.

The property of Mr. A. Grindey, and bred by Mr. Richard Terrot.

the same enthusiasm that was evinced in 1912 will be seen, and then a bright future is in store for the breed. A Club has been formed to look after the interests of the breed, and this should do much good for it.

Belgians may be approximately divided into feathered-legged birds with single combs, these being termed Barbu d'Uccle, and clean-legged ones with rose combs, known as Barbu d'Anvers. There is any amount of choice in colour, though probably the chief ones are Black, Blue, White, Barred or Cuckoo, Spangled, and Quail or Caille in the clean-legged section, and though most of these colours are found in the feathered-legged ones, the most common and popular one is undoubtedly the Millefleur, or Thousand Flowers. Rumpless ones, known as Barbu d'Grubbe, are also seen, though these, I believe, have not been taken up by our fanciers. They should certainly suit our Manx friends. The Barbu d'Grubbe are practically Rumpless Black Antwerp Bearded Bantams.

The plumage of Blacks, Blues, Cuckoos and Spangles hardly needs describing. Quails or Cailles are very pretty. In cocks the neck hackle should be of a brilliant black ground, sharply laced with buff; breast, golden buff finely laced with yellowish buff; the whole of the back to the saddle hackle golden, each feather having a black ground colour with a lacing of gold, this starting at the middle of the feather and getting narrower towards the top, forming a lance-like point, with golden silky barbs, clearer than the lacing, ending in the

upper part of the feather; the black ground colour becomes more noticeable in proportion as it approaches the end of the saddle hackle feathers; wings, the whole of the shoulders should be light gold, the lower half of each feather black, clearly defined from the upper half, which should be nankin (golden buff), shaped like a lance, with long silky barbs of a brilliant gold lustre; wing-bar, each feather should be light ochre colour, having a black triangular mark at the end, the whole of these triangles forming two regular bars across the whole width of the wing; secondaries, the bottom third of their length should be chamois colour, and the other two-thirds dull black; primaries, dull black; tail, black ground colour finely bordered with a brown line and light shafts faintly defined; saddle hackle, black ground laced with chamois colour; thighs, similar to breast; and legs and feet, slaty grey. In hens the neck hackle should be umber with a fine lacing of gold; breast, spotless nankin, the feathers towards the wings being finely bordered with a dark umber colour, and by their succession form a regular distinction to the light part of the breast; back, umber coloured feathers with a silver velvety lustre finely bordered with chamois colour; wings and



A BELGIAN BANTAM COCK.

The property of Mr. A. Grindey, and bred by Mr. Richard Terrot.

tail, very similar in character to those on the back.

The plumage of a Millefleur cock should be: neck hackle, black feathers with golden shafts broadly bordered with orange-red, each feather having a black mark at the end, terminating in a white point; breast, chamois colour, marked with a pea-shaped black spot, tipped with a triangle of white; back, red

spot on a triangle of black, these triangles continuing in equal spaces towards the tip, or end, of the wing and along the ack; secondaries, the lower third of each feather taken lengthways, which is only visible, chamois, and the other two-thirds black; primaries, black, bordered on outside with a thin line of chamois, terminating with a triangle of white. In the hen the ground colour should be an even golden chamois, each feather




A BELGIAN BANTAM.

Mr. Jamotte's Barbus d'Anvers Cockerel.

graduating into orange towards the saddle hackle; wings, the whole of the shoulder red mahogany colour, the tip of each feather white; the wing-bar russet with a black pea-shaped spot at the extremity of which should be a silvery white triangle; the dapping on the wing should be a uniform chamois, each feather at the end having a large, pea-shaped white

terminating with a black spot finishing with a triangle of white; tail feathers black, finely laced with chamois, with white tips.

In colour the Porcelain is almost unique, being a combination of straw (yellow), blue-grey, and white tips. Though a very pretty variety, it is not seen so frequently as many of the other colours.



R.I. RED BANTAM COCK.

Bred by Miss M. H. Clay.

Muffing and beard are great features in Belgian Bearded Bantams.

RHODE ISLAND REDS

Though now and again for a good few years past an exhibit of this breed has been penned, it can safely be said to be one of the latest ones to make its appearance with any degree of success. It has come rapidly to the front during the past two seasons, and now that a Club has been formed to further its interests it should make still greater progress. Splendid headway has already been made, as can be judged from the creditable specimens that have been on view during the past season. A little less size has in most cases yet to be looked for, but in type many of the birds are quite equal to those of the large variety. Another difficulty is that of colour, some being rather light and patchy, though some have been exhibited that excelled both in colour and markings, and it only seems a matter of time before we have the wee Red as good as its larger confrère.

YOKOHAMAS

This breed is another that has receded in favour, this without doubt being due to the War. Up to then it had a certain popularity,

and was seen in useful numbers at the big events. Now it is very seldom seen. The Yokohama is noted for beauty of colouring, graceful shape and stylish carriage, to which is added an abundance of lovely feathers of the finest quality, and quite a unique characteristic of the breed. The carriage and general characteristics of the birds are best described as being pheasant-like. The principal colours seem to be Gold and Silver Duckwings, Black-reds, and Whites. The colour of Duckwings is the same as in modern Game of this colour, but all black feathers must have a metallic beetle-green sheen. Lobes a bright red, and without any white, if possible. The colour of Whites, both sexes, is—beak, legs and feet, pure white or yellow; eyes, bright red; comb, face, lobes and wattles, bright red; plumage, throughout a pure snowy white, quite free from any straw-coloured tinge. Golden or Black-reds:—Cock—Head, bay; hackle, red, each feather having a black stripe in the centre; breast, thighs and fluff, black with beetle-green



WHITE YOKOHAMA BANTAM COCK.

The property of Mr. E. Brown.

lustre; back, red; saddle hackle, red, each feather having a black stripe in the centre; wing-bow, deep bay; wing-coverts or bar, black with brilliant metallic beetle-green lustre. Hen—Head, brown; hackle, orange colour, each feather having a black stripe in the centre; breast, salmon colour, lighter shade below; thighs, greyish brown; back and shoulders, brown, each feather having a light-coloured shaft; tail coverts, brown; tail, black, the two uppermost main feathers spotted or grizzled with light brown. Both sexes—Beak, horn colour; eye, ruby red; face, comb and wattles, bright red; ear-lobes, pure white (or pure red); legs and feet, yellow, willow colour or slatey blue.

Yokohama Bantams should be exactly the same in shape, colour, and general type as the large Yokohamas, the only difference being the size, which should be as small as possible, but well proportioned. According to the Poultry Club standard the weight of the cock should be 16 ozs., and the hen 12 ozs.

SCALE OF POINTS

Head	10
Quality and length of neck and saddle hackles	20
Quality and length of tail and number of feathers	25
Colour	10
Size	5
Condition, carriage, symmetry ..	25
Legs and feet	5
	100

SERIOUS DEFECTS.—Wry tail, humped back, crooked breast, more than four toes.

SUMATRA GAME

This breed has never been very popular, which is a great pity, as a good specimen is a very beautiful bird. Sumatras should be a miniature counterpart of their larger brethren, and should handle firm and muscular. The head should be small and short, with a peacomb fitting close to the head, with large bright eyes, either dark red, dark brown, or black, and full flowing black plumage, but neither soft nor fluffy, the colour of the same being of beetle-green. The carriage should be pheasant-like, straight and upright in front, and with a proud and stately action. The comb, face, wattles and ear-lobes should be black or very dark red, the gipsy face being preferred, and beak,

legs and feet, dark olive or black. Serious defects.—Other than four toes, single or rose-comb, dubbing, or any deformity.

SUSSEX.

It is not surprising, seeing how popular the large Sussex have become in recent years, that an attempt has been made to Bantamise the breed. I should say it is about seven years since Mr. F. Smalley commenced breeding the light variety, and if I am not mistaken it is about five years since they were first exhibited. In this short space of time much progress has been made in bringing the variety up to standard, though of course it is not by any means perfect yet. There are one or two breeders trying to evolve the speckled variety, and though this colour has been seen in the show-pen it will, I fancy, be a long time before this variety will claim much attention. If it can be brought to a fairly good state of perfection it should take on, for its blend of colours is a very pleasing one. The light variety, as I predicted both in *The Feathered World Year*



Photo]

[*"Feathered World."*

DUCKWING YOKOHAMA BANTAM COCK.

The property of Mr. E. Brown.



Photo] [“*Feathered World.*”

LIGHT SUSSEX BANTAM PULLET.

Bred at the Basildon Park Farms.

Book and in my column in *The Feathered World*, is becoming a favourite, and this is not to be wondered at when one considers its utility qualities as well as its exhibition ones. Type and hackle striping need a watchful eye, for these are the chief points where most improvement can be made.

The Sussex Bantam should be an exact miniature of the Sussex fowl. The general characteristics are a medium sized skull, short, strong and well-curved beak, full bright eyes, and single comb of medium size, upright, evenly serrated, and fitting closely. The neck should be of medium length with fairly full hackle, broad, deep and long body, with square breast carried well forward, with long and deep breast-bone, and a broad, flat back. The wings should be carried closely, and the tail of moderate size, carried at an angle of 45°. The legs should be rather wide apart, the thighs stout and the shanks strong and free from feathers. The carriage should be graceful and erect.

In colour the beak of the light variety should be white or horn, orange eye, comb, face, wattles and earlobes red, and legs and feet white. The plumage should be pure white, with black striped neck hackle, black in flights, and black tail, the black centre of each feather of the neck hackle to be entirely surrounded by

a white margin, and finishing in a decided white point, not a black tip or black outer edging. Serious defects are a rose comb, feather on shanks, other than four toes, or any deformity.

Though I have already pointed out that improvement can be made in type and hackle, it must not be understood that these points have not, in some birds, been brought to a good state of perfection, but usually the two points have not gone together. Some grand hackled birds are to be seen, but often enough these are not too good in shape, whilst others excelling in shape are somewhat weak in hackle. However, time will no doubt rectify these points. It is also difficult as yet, just as it is with the large breed, to get the male birds free from brassiness or stain on top, and no amount of shading appears to be able to eradicate it. Only careful breeding is likely to eliminate this.

It is an advantage to the breed that the single mating system can be used with good results, and this should go far in popularising it. In breeding choose a male bird as good as possible in hackle striping and with a clean black tail, even if he has a little smutty undercolour, but the hens to mate to him should be as sound coloured as possible. Where, however, the neck hackle is sound and pronounced both in the cock and the hens then a clear undercolour is all right.



Photo] [“*Feathered World.*”

LIGHT SUSSEX BANTAM COCK.

The property of Major J. A. Morrison, D.S.O.

OTHER BREEDS

Black Orpingtons, Langshans, Houdans, and Faverolles are now practically non-existent. A Dorking is now and again seen, but somehow this breed fails to catch on. Anconas have of late been seen with greater frequency than for many years, and some really nice specimens, both in size, colour and tipping, have been exhibited. In all points bar size these breeds should resemble the large varieties.

CHAPTER XVII

HATCHING AND FEEDING CHICKS

In the first place, the selection of a suitable broody hen is one of the most important features in hatching and rearing Bantams. It is useless putting a big hen of 5 or 7 lb. weight on Bantam eggs, for even should she be fortunate enough not to smash the lot during the period of incubation, she would in all probability crush the life out of the whole brood within a week of hatching.

Years ago I was greatly troubled over sitting hens breaking eggs and trampling chicks to death, so I determined to remedy the grievance by rearing suitable hens for hatching Bantam eggs.

What I did was to purchase three or four Buff Pekin hens, and these I mated to a black-faced White Silky cock, and from this pen I hatched some twenty or thirty pullets, which for sitters and mothers were worth their weight in gold. Sit! why, they would sit in the middle of the road on a few small stones; or, if you are hard up for room, you can put them on the top shelf in the scullery, or on the mantelpiece, and they will sit from three weeks to three months if required. They will sit anywhere, and, as a rule, will rear two or three broods of chicks each year, for I have never known them lay more than a dozen eggs before becoming broody, and are so quiet and reliable that no Bantam fancier should be without them.

Having secured a suitable hen, which should not weigh more than 3 lb. at most, before putting her on the nest, give her a good sprinkling with insect powder, and see that she is perfectly healthy and, if possible, not scaly-legged. The latter point is of great importance, especially where smooth, clean-legged chicks are a necessity, as in Game Bantams. I fancy I hear many of my readers say, "But what effect can a scaly-legged broody hen have on the chicks she rears?" My

answer is: A very marked effect, and if you have never had scaly-legged chicks before, you can depend upon having them in the first brood you rear under a scaly-legged hen. This is a wrinkle, and you would do well to bear it in mind.

Now as regards nests. I very rarely use anything except the usual orange boxes, which can be purchased from any of the greengrocers for a small sum. I lay these down on their side, and nail a strip of wood about four inches deep along the front. You will then have three good roomy nests. The next thing to do is to get plenty of nice soft hay, and make a nest in the shape of a large basin, i.e. hollow in the middle and nicely rounded at the sides, to prevent the eggs from rolling out when the hen leaves the nest to partake of food. There is an old saying, which many breeders follow, I have no doubt—that is, before putting in the hay, they will tell you it is necessary to have soil at the bottom. This is all "bunkum." And, again, they will tell you that they must be set on the ground. Let me tell you that it is quite unnecessary in both cases.

Having put the hen on the nest, give her a dummy egg or two until she properly settles down, which should not take more than a few hours. If at the end of that time she does not attempt to fly off when you place your hand under her, it will be quite safe to put the proper eggs under, and this is best done at night time. Early in the season, when cold winds and frosty nights are prevalent, it is unwise to give a hen her full complement of eggs; better two less than one too many, for if she has too many it will make her uneasy, and in all probability she will spoil several eggs in her endeavours to get them all properly under. At the end of four days after setting hens, examine the eggs by holding them before a candle. This must be done at night-time; the clear eggs can then be taken away, and are easily distinguished from the fertile ones, the latter looking cloudy, and the veins of the germ will be quite distinct. By taking out the unfertile ones, in addition to making more room for the fertile ones, it gives them a much better chance of keeping the proper temperature.


The front of the nest should be covered either by a piece of board or sack, and each night carefully remove each hen to feed, and keep her off for ten to fifteen minutes. The best food for sitting hens is maize, and let them have plenty of water and a good supply of cinder ashes for them to dust in. Should the nest become fouled or any of the eggs

broken, take out the nest, replace with fresh hay, and wash the eggs with lukewarm water. If the nest is properly made, nice and soft, and the hen allowed off sufficient time each night until the wants of nature have been attended to, and providing she is a suitable hen, as I have already described, this trouble will very rarely occur.

At the end of nineteen days the eggs will be hatched, if they were fresh when put under, and not later than twenty days at the most. If you would have strong, healthy chicks use fresh eggs, and the larger the better.

his brief sojourn in this world will be a misery to himself and an eyesore to those around him.

Returning to the sitting hen, on the eighteenth day of incubation, when she comes off the nest for her usual feed, dust her again with insect powder, and do not disturb her again until the next night, when the chicks ought to have put in an appearance. Let hen and chicks remain quiet until the next morning, when you can give them their first feed. Before doing so, give the hen a good feed of maize and water to drink, or she will be liable to eat all the food away from the chicks. Nature has so provided



[Photo]

ANCONA BANTAMS.

[A. Rice.

Father, mother and daughter—the property of Mr. R. W. Tunstall.

FEEDING CHICKS

Do not attempt to hatch and rear Bantam chicks by artificial means, for you cannot do with any degree of success. Incubators and foster-mothers are all right for the larger breeds of fowls. I do not say that you cannot hatch Bantam eggs by incubators, as with a good reliable machine and careful attention daily, and by allowing the egg-drawer to be open from fifteen to thirty minutes daily, according to the period of incubation, you can obtain very fair results, but the Bantam chick, hatched by artificial incubation, will invariably be a very fragile, puny little mite, that will be quite an impossibility to rear, and during

that the chick does not require food for at least sixteen to twenty hours after hatching, whilst in the large breeds twenty-four hours can elapse before giving food.

The first feed, then, may be egg boiled for about five or six minutes, just long enough to cause it to be set; this should be chopped up fine and mixed with equal parts of coarse oatmeal and stale bread crumbs. Continue this feed for the first two days, by which time the brood should be well on their legs. On the third morning give them for the first meal Spratt's chicken or game meal scalded, and allowed to stand until nearly cold. The next feed should be given about two hours later,

and this should consist of pure oatmeal mixed with sweet milk until crumbly, or stale bread crusts soaked in water and mixed with oatmeal till crumbly. This should be given at least three times a day, with the addition at noon-day of a little lean meat, chopped fine, for the first two months. When the chicks are a fortnight old, groats can be given for the last feed at night, and when two months old wheat should be substituted.

The question has often been asked how frequently should chicks be fed? To a great extent this depends upon the breeds, but for Bantams every two hours for the first week, and from that time till they are three months old, four times daily will be sufficient. From three to five months, three times a day. A fair allowance of animal matter, such as lean meat chopped fine, granulated meat, or maggots, should be given to chicks once daily, especially during February, March and April; this will enable them to withstand the cold winds and damp. Where chicks have not the liberty of a grass run, boiled vegetables mixed with the soft food should be given at the noon feed. Above all, feed regularly.

We now come to the water question, and which has been freely discussed in *The Feathered World* from time to time. My advice to you is the same as *Punch's* on the marriage question: "Don't." I know of many successful breeders and exhibitors who rear from 200 to 500 chicks every season, and the only water they get is what they can obtain from the grass after a shower or heavy dew. For my own part, I do not believe in water for chicks. Since I discontinued the use of water for my chicks I have not had one solitary case of diarrhoea, but let them have access to water, especially in hot weather, then your trouble begins. If you never give chicks water they never require it, and are much healthier without; but once you start giving them water, you must continue till the end of the chapter. If your chicks have a grass run and are fed as stated above, they will not require water. Of course, when the bird reaches the age of, say, four months, when grain will necessarily be his principal food, then, and then only, need water be given.

When chicks are old enough to be taken away from the hen—viz. about three months—the cockerels should be separated from the pullets and put on different runs where they cannot get together. By doing this you will find that both sexes will thrive better and

settle down more contentedly; and a good plan is to put an old cock along with the cockerels to prevent their fighting and to maintain order.

Although chickens do not require water, the hen must not be forgotten, and to meet the necessity I fasten one of the ordinary drinking tins inside the coop, about eight inches from the floor. This enables the hen to drink what she requires, and at the same time it is out of reach of the chicks. The tin is easily kept in its place by a couple of wire staples.

During February, March and April I prefer false bottoms to the chicken coops, on account of the damp, cold weather generally experienced at that time of the year, and by having them loose they can be very readily removed for cleaning. When the warm weather sets in it is advisable to take out the bottom and allow the chicks to sleep on the ground; but where rats and other vermin abound, the bottoms should be kept in all through the year. It is advisable also to have a hinged shutter to the coop for protection at night, and which also affords a shade from the sun in the daytime; the shutter can be lowered to any desired height by having a chain attached. The coops should be limewashed every five or six weeks; this is very important, if you would have strong, healthy chicks.

FEEDING ADULTS

In the notes above I have gone fully into the feeding of Bantam chicks, but it is necessary to carry the subject somewhat further and throw out suggestions for treatment as the adult stage is reached. For unless all kinds of Bantams are judiciously fed and cared for it will result, to use a Shakespearean expression, in "love's labour's lost."

A Bantam so brought up that size and feather are ignored is a Bantam ruined. Of course, it is possible that failure in these respects may be the result of breeding, but it very often happens that from the very best breeding pens procurable, the stock, through being kept on wrong lines, result in failure.

I have seen Game Bantam chicks, that have been given a liberal diet of cooked meat, scraps etc., and other soft food, in addition to a plentiful supply of grubs and worms that the chicks may have picked up in the kitchen garden, grown so large and feathery that by the time they reached four months they were much too large and coarse even for adult birds, larger even than the parents they were bred from, consequently useless either for exhibition

or breeding purposes, and simply because they had been brought up on unsuitable diet.

The above diet is all well and good for pushing on big breeds where size and bone are required, but in Bantams the order is reversed. Feeding, therefore, becomes a matter for the highest consideration. In Bantams size must be kept down, and discretion used in feeding for hard or soft feather. As a general principle it may be set down that hard feed produces hard and short feather, and vice versa. ¶



[Photo]

[*"Feathered World."*]

WHITE FRIZZLE BANTAM COCK.

Bred and exhibited by Mr. A. Hobson.

Breeders of Game, Malays, etc., will therefore proceed on the lines of hard corn as staple food, with only small additions of soft food, the less the better; whereas breeders of Rose-combs, Pekins, and kindred soft-feathered stock will judiciously place soft food first, and give hard corn merely as a nightcap before roosting time. In neither case will the food be chosen of too fattening and frame-making a nature, yet sufficiently nourishing to build up and keep up a sound constitution. For if the birds be brought up in too starved a condition, weakly debilitated frames will result, and lead to much after trouble

Where birds have full liberty on a good grass run less food is required than if they are kept rigidly confined to a pen. In all cases a little soft food should be given in the morning, composed of Spratt's Biscuit Meal scalded with hot water, and brought to a fine crumbly state with fine sharps, or what is sometimes called "thirds"—a meal much resembling flour. A very little maize meal may be added. One handful of this mixture to two or even three full-grown Bantams will be sufficient for morning feed, presuming the birds have full liberty; at night, before roosting time, good sound wheat in a similar proportion. A change to *dari* occasionally will be useful, and be greatly appreciated by the Bantams, for, like their bigger brethren, they like change of diet. In the early spring hempseed should take the place of *dari*, as it furnishes natural heat, and will be found very beneficial in the production of eggs, especially where the birds are kept in exposed positions. Buckwheat is also an excellent food, but, as a rule, birds do not take to it and will rarely eat it unless given separately from other grain. I do not advocate mixing the grains; this precludes the possibility of change. Where birds are kept in confined runs, the same menu may be given, but it will be well to add any available household scraps, such as boiled cabbage, bits of meat, etc., chopped fine, avoiding a too free use of boiled potatoes, or their peelings, which are most fat-forming, therefore unsuitable for Bantams.

Never stint the supply of green food; young dandelion, watercress, lettuce, young cabbage, in fact almost anything that is green, should be given to closely confined birds daily, whilst the cooked meat chopped fine given three times a week will be sufficient. If the green food be introduced largely in the pen, it should be removed at once as soon as its freshness has passed away. Nothing can be more harmful than decaying vegetable matter. And with regard to both corn and meal, see that it is not damp or mouldy, for nothing I know of will set up disease sooner than this. Wheat especially is injurious, if it gets into this state. All food stuffs should be kept in large airtight tin canisters or bins, and in this way kept secure from damp and mildew. The flesh meat, too, should be thoroughly wholesome. Some people seem to have an idea that any sour, putrid stuff will do. This is a grave mistake. I find Brand's Meat Meal, a lean meat fibre, very beneficial, both in rearing chickens and given

in small quantities to adults in morning meal during February, March and April, when eggs are at a premium. Spratt's Crissel is also an excellent addition to the soft food, and I have heard it highly spoken of. I do not advocate poultry spices. They will doubtless produce eggs, but at a great cost of constitution, and a Bantam has not enough original vigour to stand such drastic helps. Where birds are confined the grit supply is very important; shell and flint mixed, which can be had from 5s. per cwt., will well pay for its cost. It should be given in a box, and not scattered about anyhow in the run. The only thing that will bear scattering is the corn. If this can be flung amongst loose straw, bracken, chaff and the like, the search will always afford useful exercise for the birds.

The soft food is best given in wooden troughs. These can be thoroughly scalded out from time to time. Never leave any food to be soured by the sun, nor yet to form an inducement to vermin. In confined runs a few shady boughs of trees should be laid over the runs, or some kind of awning be put up in tropical weather, not only to keep green and other food fresh the longer, but to furnish a welcome shade for the birds.

It will be found that eggs from Bantams come much more freely, as in fact they do from any breed of fowl, in warm weather than they do in the cold spring months.

If eggs are wanted early, see that the soft morning food is given hot, as reasonably hot as the bird can take it, and that they are not fed in the evening so long before roosting time that their crops have got rid of the corn before retiring, and so are left to pass the cold frosty nights without the natural essential for keeping them warm.

WATER

Water, which we deprecate for chicks, should not be withheld from adult birds. In fact it is absolutely necessary for them. It should be given fresh twice each day; if the weather is sultry three times or even more. Under all circumstances keep it as much out of the sun as possible. Use earthenware vessels well rinsed out each time. Studiously avoid the use of zinc or lead vessels.

As a tonic and for keeping the birds in good health, I put a small piece of sulphate of iron about the size of a small bean in the water twice a week. Twopennyworth will last months. This must not be put into zinc drinking tins.

I think I have gone fully into all the details of feeding and the proper food to be given, and will now pass on to

HOUSING

The Bantam fancier who flatters himself that any sort of a dirty, damp, draughty place is good enough to bring up and keep his pets in, cannot possibly make a greater mistake, and so first-class men who know their business never attempt it. Housing correctly is quite as much a panacea to healthy condition, whether for breeding or exhibition, as correct feeding. In fact, to put the matter shortly, all the work of mating, rearing, feeding, etc., may easily be rendered abortive by a week's housing in unsuitable quarters, and such a fiasco seems deplorable in the face of the fact that Bantam quarters, good and suitable in every way, can be had nowadays at a most reasonable figure; in fact, no kind of poultry house or pen costs so little as those for our pigmy pets. First of all, the place must be weatherproof. So many of us are amateur carpenters, not a few think they can construct a Bantam house for themselves. And, we ask, why not? I have seen almost as good houses turned out by the amateur as by the professional carpenter. But in amateur carpentry the temptation comes in to use any kinds of odd timber that is lying about, odd lengths, short and long, and so it very often happens that when all is put together there are cracks and chinks showing here and there when we view our work from the inside.

Now this must be avoided. Draughts must not be allowed; better far let the birds roost out in the open than exposed to a draught. The wind blowing in from a hole the size of a Bantam's quill, on a cold stormy night, direct upon the face of a bird, will leave its mark in the shape of a swollen face, bad cold, and sometimes roup.


Stop all such crannies with putty or wooden plugs. A home-made house is all the better for being covered with felting. Get the best. It is cheapest in the long run. But if our Bantam house has to be draught-proof it must not in consequence go unventilated. This is most important. More cases of roup are caused every season by allowing birds to be cooped up in badly ventilated houses than from all other causes combined. Birds that are allowed to roost in large numbers huddled together in evil-smelling, badly-ventilated places all night, come out into the cold morning air more dead

than alive, and the contrast of the sleeping quarters and the outside atmosphere is so great that the birds catch cold and are soon down with roup and kindred ailments.

The house should have a current of pure air passing through from end to end, well above the birds' heads, the higher the better. These ventilation holes are best if provided with shutters that will let down and take up, and even off. Two kinds should be to hand, the one of solid wood in case of very high, cold

happily, I think these exhibitions are few and far between.

Game Bantams are by nature very pugnacious, consequently dubbing is a necessity to prevent them tearing each other's wattles and combs to pieces. The misery caused by two undubbed birds getting together and having a pitched battle is greater far than is ever experienced by the operation of dubbing, which should never occupy more than from three to four minutes at the outside. Those



LEAN-TO BANTAM HOUSE AND RUN.

winds in winter, the other a frame covered with perforated zinc for cold gusty weather in general. In fact, it is a good plan to so have the frame in which they slide constructed that both can be used together. In this way just the requisite amount of fresh air could be given that circumstances called for.

DUBBING

Much has been said on this vexed question both for and against, and at one or two of our shows dubbed birds are still prohibited. Now,


people who have been trying to put a stop to dubbing have never dubbed a bird in their lives or yet seen one done. I have dubbed some thousands in my time and have never yet lost one single bird either under the operation, or from the effects; but I have lost many a valuable young bird through fighting before he was dubbed.

If the young fancier who reads these notes has never performed the operation, and he wishes to do so, he should first of all get a pair of proper dubbing scissors. These scissors are

curved in the blade in order to fit close to the head and enable the operator to get a clean cut. An assistant should hold the bird firmly, taking one leg and one wing in each hand, and should stand facing the operator. Nothing else is required except a bowl of cold water and a sponge. The operator should then take hold of the bird's comb with the forefinger and thumb of the left hand, and hold the head firmly. The ear-lobes should then be removed,

his run. In the majority of cases you will find that as soon as ever you put him down on the ground after the operation, he will at once commence crowing. What better proof than this can you have, that the operation is not by any means so painful as some would try and make it out to be.

The day following the operation, a little vaseline should be applied to the sores. This will help nature in the healing. Nothing



BANTAM HOUSE AND RUN.

then the wattles, cutting the latter from back to front by one clean cut. He should then insert his thumb inside the beak of the bird, and grip the beak with the thumb and finger, stretching the neck to the full length. Then, pressing the scissors as close as possible to the head, cut off the comb with one long clean cut from back of the head to the beak. The head should then be bathed with the cold water, which will at once stop any bleeding, and if the weather be warm the bird can be returned to

further is required, and within ten days all traces of the operation will have disappeared.

HOUSES AND RUNS

With regard to size of Bantam houses, all will depend upon the number of birds kept together. For a pen of half a dozen hens and a cock, the house should measure about 4 ft. by 3 ft. From the floor to the eaves should be at least 2 ft. It would be better 2 ft. 6 in., allowing 6 in. more to the ridge of roof. Under

the floor should be a shelter reaching 18 in. from the ground, and boarded all round with the exception of the front. This shelter will prove extremely useful to the birds in bad weather. Into this shelter put a good supply at frequent intervals of peat moss, chaff, or cut straw. This will keep them dry and comfortable in the time of cold winds and wet weather. Another capital plan, if the top of the run will bear it, is to cover it with a layer of bracken and fir branches during the winter months. It is wonderful how such an arrangement keeps the snow off. Melting snow has a very bad effect upon Bantams' feet. It is pretty generally known that any solid substance like ice or snow when melting absorbs surrounding heat, and with the absorption of heat from the foot, chilblains, etc., arise, the feet will become blistered, and in some cases the toes even will be frozen completely off.

On the floor of the run it is best to have fine dry sand, such as sea-sand or very fine gravel, and kept fresh and sweet by removing the droppings and raking the sand over once or twice a week. A dust bath must also be put into the shelter, otherwise your pets will soon be swarming with vermin. The best dust bath is fine ashes to which a small quantity of sulphur has been added. This should be put into a small box about 18 ins. long and 6 ins. deep, a fresh supply of ashes and sulphur to be given at least once a month. Where possible, the run should be on sloping ground, so that the water is naturally drained off it. During the winter the runs should be boarded quite 18 ins. or 2 ft. from the ground. Shelter like this is absolutely necessary if the fancier's object be early eggs, and early eggs mean early chicks in readiness for the early chicken shows. Not only is it a case of early eggs, but of a better egg production all through.

Keep plenty of moss litter or sawdust in the house itself, and never adopt the merely occasional method of clearing out. It may answer all right for cross-bred laying fowls, but Bantams are not built on the same lines. See that your houses are well done once a fortnight at least. Give them a good lime-wash from time to time. Into the lime-wash add a little carbolic acid. This will effectually kill fleas, ticks, etc. Clean the floor well. Put fresh litter or sawdust down, to which has been added some disinfectant. Then if ventilation be properly attended to you may expect your birds will keep in good health. Remember to put up proper perches. Eschew all square

perches or flat boards unless you wish to ruin the birds' feet. The perches for all Bantams, with the exception of the feather-legged varieties, should be round and about the thickness of a broom handle, made so that they can be easily taken down and cleaned.

Houses such as I have described, and which I have found to act admirably, should be of three-quarter inch boards, tongued and grooved, with door and window complete, and it is, therefore, the height of folly, when valuable first-class stock has to be kept, to put up with any kind of a make-shift house, and run the risk of disease and probably ultimate loss. It will be found that the initial expense is after all the least expense, for once get birds thoroughly out of health, it is quite a question whether they will ever again be really fit for either exhibition or breeding.

Whilst on the question of houses, let me warn the young fancier against the unscrupulous advertisers of cheap poultry houses. Thanks to the Editress of *The Feathered World*, when these fraudulent transactions are made known to her, the columns of *The Feathered World* are closed against them once and for ever and this is as it should be.

Before leaving the subject of housing, see that they are free from all access of vermin. A solitary weasel in half an hour may do £100 worth of harm, and rats are by no means particular if they cannot find grain handily, they will go for small birds like Bantams. The boarding of the run is a great protection against vermin, and if the mesh of the wire netting immediately above the boarding be small, this will act as a further safeguard. It is well to have a wire covering over the run, too, especially if it be in the neighbourhood of a wood. Some of the larger kinds of Hawks, Carrion Crows, and Ravens have been known to swoop down upon small Bantams and carry them off. A covered-in run is a safeguard, too, against dogs and cats. Stray dogs and cats are at times a source of uneasiness to every keeper of dwarf poultry; but wire netting is effectual against them, and prevention is better than cure. Attention to such details as I have here set forth on the part of Bantam fanciers will well repay them. It will be found on trial that the old adage so often quoted will come true in the case of Bantams as in that of most other things—"A thing that is worth doing at all is worth doing well."

WASHING

Many a good bird has lost his position in the prize list for the want of a good wash, and others have had the same fate through being badly washed. Therefore it behoves the young fancier to pay the strictest attention to washing birds for exhibition.

All light-coloured birds, such as Piles, Game, Duckwing, White and Buff Pekins, White Rosecombs, Frizzles, Polish and Booted require to be frequently washed and properly dried.

In the first place then three bowls of water are necessary, a large bath-sponge, an ordinary

of your penknife, or, better still, with a match-stalk sharpened to a fine point, which can be more readily inserted under the scales, and all dirt taken out. Having done this, the legs should be again scrubbed with soap and water.

The bird should now be held in the water with his head just above the water. Allow him to remain in this position a few minutes to thoroughly soak the feathers, and having done so, the bird should be allowed to stand up, and the soap should be well rubbed into the feathers, rubbing the feathers well all the time with your hand from head to tail and



WASHING A BIRD.

nail-brush, a bar of good soap (Sunlight or White Curd Soap for preference) and a plentiful supply of *soft* water; never wash in hard water when you can possibly avoid it. Bowl No. 1 should contain warm water, so warm that you can comfortably bear the back of your hand in, and sufficient in which to immerse the bird, just leaving the head above water when held down in the bath. The comb, face, and wattles should be well scrubbed with soap, using the ordinary nail-brush for the purpose, then wiped dry with the sponge. The bird's legs should then be well soaked in the water, holding them there for three or four minutes. Then take the nail-brush and soap, and give them a vigorous scrubbing, and if there is any dirt left underneath the scales, this should be removed with the point

squeezing out the dirt. Repeat the operation until all trace of dirt is removed. Immerse the bird in the water again, to get out all the soap you can, then place it in water No. 2, which should also be warm, but not so warm as No. 1. With the sponge rub the clean water well into the feathers, and this must be done thoroughly, or all your labour will be in vain, for if any trace of soap is allowed to remain the bird will be in a worse condition at the finish than when you commenced. The bird can now be taken out and put into bath No. 3, which should contain water the warmth of new milk, or just with the chill off. In the case of white birds a little blue should be added to the third water, just sufficient to tinge the water slightly, but be very careful that you do not put in too much. In washing

Barred Rock Bantams and Scotch Greys, more blue must be added, and this will improve the ground colour considerably. In using the blue, it should be placed in a piece of flannel and tied up, put in the water, and as much squeezed out as the case requires. In washing buff birds, of course, no blue whatever is required.

After having sponged the bird thoroughly in the third water, all the water possible should be absorbed by the aid of the sponge, and if the sponge be a large one of good quality, the bird can be half dried by this process. Then remove the bird from the water, and wipe him as dry as possible with the aid of a towel, after which it should be placed in an unlined exhibition hamper or one of the ordinary exhibition wire pens (the former preferred), with some nice clean hay or straw in the bottom, and the hamper should be placed about two feet from a nice bright fire, and some kind of screen placed at the back, such as a newspaper or piece of canvas, to keep off the draught. The bird should not be allowed to stand in one position too long, but should be made to turn round every few minutes, in order that the feathers may dry equally all over. This should be continued until he is almost dry, when he will begin to dress the feathers himself. The following day the bird should be well rubbed down with a silk pocket handkerchief, in order to web out the feathers properly.

It is always advisable to wash birds at least three or four days before they are required for exhibition, for if a bird is washed one day, and sent to the show the next, it is very liable to contract a cold on the journey and also the feathers do not have time to get into proper order.

I should advise novices to practise on common birds first to get used to the process.

TRAINING AND PREPARING FOR EXHIBITION

It is necessary in a book of this kind to say something upon the way Bantams should be trained and prepared for exhibition. Apart from the necessary washing of the white and light-feathered birds, there is a certain amount of schooling through which all fowls, *but especially Bantams*, should go, if they are to make their successful début in the show-pen. Fancy a lad tumbled into life without any previous preparation for that which he was to encounter, raw, crude, and unsophisticated. Should we expect him to win all along the line?

Even so is it with our feathered pets. They are unsophisticated in exhibition ways, and have to take their lessons like other things. Many a good bird has been quite spoilt through lack of training. Certain things are required of them in the show-pen; they do not do them, and the judge has neither time nor patience to stand waiting for them whilst they pick up by natural intuition, if they can, what should have been taught them before they came to the show. For instance, the bird will not settle, runs about its pen in a wild, frightened manner, sticks its head in a corner, and, if disturbed, dashes against the sides and top of a pen to the disgust of the judge and no small injury to itself at times. In the end it is passed over, naturally so, and marked in the judging book "wild in pen." Now, a very little care would obviate this, if it were exercised at the right time. If, when the bird was some four or five months old, a few minutes' training in a pen had been gone through daily for a short time it would have removed all chance of a fiasco when exhibition time came round. What I do myself is something after this manner: I generally take the bird off its run in the evening when it has gone to roost and put it into a suitable training-pen for the night. Such a pen is about 18 ins. by 18 ins., and can be procured from any penning firm for 1s. 9d. each. They are clean, light, and airy, better by far than home-made wooden ones, and are, in fact, such as the bird will have to be exhibited in. Here, in the pen, I leave him for a quiet hour or two to settle down a bit; then, when it is quite dark, I take a candle, and go and stroke him *very quietly* from the back to the tail, then from the throat to the thighs, talking kindly to him all the while, and touching him very lightly and gently at first.

It is astonishing how much a bird will stand by artificial light that it would not endure by open daylight. When I have handled my bird till I have succeeded in getting him to stand still in one position, I take a small stick and repeat the operation with that. If the bird is inclined to carry his tail high, I repeatedly pass the stick over his back and tail, gently pressing the tail down as I go, to induce him to carry it lower. This habituates the bird to the feel of the wand on his back, and he gradually associates with that feeling the passing of the wand over the tail. I have known birds well trained instantly to lower their tail the moment the stick rested upon their

back. They acquire, too, a habit of slightly walking away from the stick, and naturally lower the tail to be free of obstacles. If he droops his wings, which they often will, then I touch him under the wings, lift them up, and in every way try to induce him to shrink himself into proper shape and form before I leave him. Even very good birds have sometimes naturally more or less bad habits of wings, tail, etc., etc., but they can—if there is really nothing radically wrong with them, and it is merely careless habit acquired in their free state of chickenhood—be easily schooled out of them. A few gentle taps on the back of the legs or toes will greatly improve the style and carriage of a bird. The training having progressed thus far, I next secure some tit-bits, bread sop, or a little meat, and hold it high up in front of the pen, to induce the bird to reach up, a great desideratum in the Game Bantams, but in the soft-feathered varieties it is a thing of lesser, and, in some cases, of no moment. This kind of training is carried on for a few nights. The bird may then be approached in the daylight, and more and more familiarities indulged in till finally such a stage is reached that, when he sees you coming, he gets himself pretty well prepared beforehand, knowing well what *is coming*. Birds that have been handled very much in their chickenhood require very little training indeed. Sometimes after the chicks have left the old hen, for safety we have had to remove them by hand every night to their proper roosting places, which they could not reach of themselves from the runs. Such birds have most easily become trained and ready for exhibition; in fact, we have known such successful at the show when they have hardly been in the training-pen half a dozen hours altogether. A little bread and milk sop, with a little lean fresh meat cut up fine and given to the birds for a week before the show, will invariably get them into good show condition. A little canary seed and millet, too, is very beneficial if given once a day through the training period. Like a horse that has been well broken, so a bird that has been well trained never forgets the discipline he has been through, and is well-mannered on all subsequent occasions, the pride of his breeder, and a delight to the judge, if, added to a thorough training, he is also a first-class exhibition specimen.

But there is a little more yet to be done, even with a well-trained bird, before he reaches the show-pen. He must be suitably despatched

to the show. Most of our best shows are in winter, and it does not do to send birds to exhibitions in flimsy structures, ill-lined, and at the mercy of any great box or parcel to crush them. See that you have good, strong, sound baskets, lined with comfortable material, and if the weather be very severe a double lining will be better still. I have known exhibitors go one step further, and sew sacking round the outside and over the lid of a double-lined hamper. If many birds are going to the show, it is better to have partition-baskets. In this way the birds are kept warmer. Each tends to heat the basket, and they mutually keep each other warmer than if they went in separate baskets, then it is cheaper. Four or five birds can travel in one basket at about the rate of two separate baskets. There is not so much risk either of the baskets going astray. In every way, both on the score of convenience and economy, this is the best way to send birds to the exhibition. See that each compartment is duly padded well with dry hay at the bottom to prevent the birds suffering a severe shaking when the baskets are roughly thrown down at the stations. Some porters have anything but a gentle way with them; they seem to take a delight in smacking down hampers as if they were cast-iron.

Before starting the bird off to the show, give it a feed of milk sop. Soak the bread in cold water first, squeeze out all the water with the hands, and add good fresh milk. Remember that anything of a sour nature tends to give the birds diarrhœa, and unfits them for show. On the return of the bird it is necessary to give the same food, no matter whether it is 1st and special or only vhc. It must be well cared for or it will feel the effects more and more of both journey and exposure in the show.

Keep the bird isolated for a day or two on its return as a precaution. One can never tell what disease a bird may have contracted at a show, and it is possible that you may thus contract in your yard something highly undesirable, such as roup or diphtheria, unless due caution is taken. If the bird be kept by itself for a few days, then you will be enabled to see whether it is all right before turning it down with the rest. Nothing is more common than for birds to return from show with some contagious disease, generally contracted from the drinking vessels, which have been provided in the show-pens, but which nevertheless have, in many cases, never been properly washed

out or disinfected after a previous show. This is a point that show officials and pen owners should be extremely careful about. The wonder is that public attention has not long since been drawn to the matter. If the cups after a show were turned down into a solution of Jeyes' or Condy's Fluid, the risk of contagion would be much minimised. The diseases contracted from foul cups are unfortunately of the worst and most virulent type, and the harm done in yards from such sources is incalculable. Only very recently some birds, under four months old, were sent to a show; they won. They were on their return put with the rest. In about a week, though previously there had not been a vestige of roup about the place, almost every bird in the yard was going about with swollen faces and closed eyes. Fortunately it was promptly dealt with, and eradicated about as quickly as it had arrived. But in the case of real roup and diphtheritic roup, it is unfortunately a thing that cannot be so summarily dealt with, so that every care should be taken by those who supply cups at exhibitions to see that they are clean and disinfected. There should be a special cleansing after each show. By this means, we should soon hear less and less of these hateful diseases, and our birds would be a less cause of anxiety to us.

CHAPTER XVIII

DISEASES

WE now come to the final chapter of our work, and close these hints and instructions on what is to many people a most fascinating hobby by some reference to the diseases to which our favourites are liable, and from which we may in a measure relieve them. Bantams, being so much in-bred, are consequently weaker in constitution and more liable to certain diseases than their bigger brethren, though, of course, there are others from which they hardly ever suffer, more particularly leg and foot weakness, so prevalent in the heavier breeds. Bumble-foot and hock trouble seldom or never attack Bantams owing to their slender build, but, on the other hand, we find Bantams more liable to egg trouble, owing to their diminutive size and, in Game Bantams, their peculiar shape, which requires them to be bred too fine to always allow of the free passage of

the egg, so that this quite makes up for their immunity from such diseases as before mentioned. Everyone feels it hard to lose a good bird from disease, and especially if it has been a heavy drain on the exchequer; still, it seldom pays to doctor fowls, our experience being that in almost every case a bird that has passed through a very severe illness becomes useless either for showing or breeding from. We like to do our best for the suffering one, but often it would be wiser to put the patient at once out of its misery, both for its own sake and that of its owner. Slight ailments are, however, quite a different matter, and may be dealt with successfully.

Roup is, perhaps, the most prevalent of all diseases, and is nothing more or less than a neglected cold, which, if treated promptly, can be nipped in the bud. A free dose of salts, as much as would lie on a shilling, keeping the birds in a warm room free from draughts, and feeding on warm bread and milk, will do more to cure than all the patent pills in the pharmacopœia. I am not a believer in pills. The more you give a bird the more it seems to want. However, for those who do pin their faith to all and everything advertised, they will find their wants supplied freely by application to "specialists." A friend of mine, however, who is somewhat susceptible to chills on slight occasions, is a great believer in half a dozen drops of terebene on a lump of sugar when he feels the cold coming on, and this has led him to try the same remedy (on the first discovery of cold) upon his Bantams with good effect. Occasionally he varies it with eucalyptus oil, or mixes the two, and if the bird shows signs of emaciation, administers about four drops of the compound in cod-liver oil. The results are, he says, invariably successful. Still, keep your birds in good, well-ventilated houses, give them plenty of fresh water as often as possible, with a bit of sulphate of iron as large as a pea dropped into it, feed on good sound dry corn, and let it be given them as advised in previous chapters, and you will be little troubled with roup. If, however, your birds contract roup at a show, treat as described as soon as you notice any watery discharge from the nostrils or eyes. If taken in time before the thick yellow, disagreeable smelling deposit comes on the beak, four or five days will, in the generality of cases, effect a cure.

Diphtheria is the most contagious and difficult of all diseases to cure, and though I have cured dozens, still I think the safest

plan is at once to kill the bird and burn the body, as by trying to cure *one* you may spread the disease through the whole yard, by carrying it in your clothes, feeding vessels, water pans, etc. In this way you may very easily lose your whole stock, and bitter as may be the first pill, especially if the bird affected is a valuable one, it is perhaps, after all, the least difficult to swallow. Having taken the drastic method in hand, next proceed to limewash all the places which the bird has used for roosting in, etc. Add a few drops of carbolic acid to the mixture, or, if this is not available, try an ounce of permanganate of potash. You will get a purple colour instead of the pure lime-white; but, never mind, you have secured a powerful disinfectant, and that is really what you want. With the disease checked, take care not to let your birds contract it again, and a safeguard will be found against this in the sulphate of iron in water, as before recommended.

Cholera is another disease difficult of cure. It is also highly contagious. The symptoms are diarrhoea with excessive thirst. Simple diarrhoea can generally be checked with a good peppering of prepared chalk over the food, which for a day or two should be rice boiled in milk, but when the more aggravated symptoms of cholera put in an appearance, it is better to put the bird down at once. Thoroughly cleanse all feeding troughs and drinking vessels, and limewash as before. A little camphor and a drop or two of laudanum in the water will prevent the disease spreading.

Cropbound is generally caused by foreign matter of a stringy nature being taken into the stomach, such as horse hair, dried grass or straw, and which obstructs the outlet of the crop, causing the crop to become quite hard. The best plan is to fill the bird's crop with water syringed into it as hot as it can bear it. Then carefully knead the hard matter with the hand till it shows signs of dissolution. When it seems to have assumed a fluid state, eject the same by turning the bird's head downwards, squeezing all matter out of the crop with the hand. If this process be repeated about three times it is not difficult generally to wholly remove the offensive matter.

The bird will then require feeding on soft food, milk sop for preference, for a couple of days. No hard corn should be given for at least two days.

Slight cases of crop-binding can easily be treated by confining the patient to a small pen, allowing it *plenty of water to drink*, but no

food for two clear days. If at the end of that time the contents of the crop have not diminished at all, it will be necessary to operate by opening the crop. This requires great care or the result may prove fatal; although I have operated on scores of cases with most satisfactory results.

In the first place then, an incision should be made, choosing a place near the top of the crop, free from any large blood-vessels, and after plucking a few feathers so as to give a clear ground, make a cut about $1\frac{1}{2}$ ins. long. Take the handle of a *small* teaspoon, and by this remove the contents of the crop, afterwards oil the little finger well and insert inside the crop, passing it round the crop to see that nothing is left, and that the outlet is quite clear. After this has been done wash out the crop with warm water to which a few drops of Condry's Fluid has been added. The edges of the wound should then be stitched together with white silk thread. Great care should be taken to see that both the inner skin and outside skin of the crop are stitched separately and *not together*. After sewing up the edges of the wound a little vaseline should be applied; nothing more is then required. The patient should be fed on bread and milk, not too moist, for the next thirty-six hours, and neither grain nor water should be given during that time.

Gapes is a trouble which affects little chickens in some localities more so than in others. This is caused by a thread-like worm lodged in the lining of the throat. How it comes there is more or less a matter of conjecture; still, when once it is there, if it be not removed or killed, the chick will speedily lose its life. The best plan is to place all the affected birds in a small box and treat them to a thorough dusting with camlin. This should be blown into the box through a small hole by means of a small bellows supplied for the purpose. This powder incites coughing on the part of the chick, and in the act the worm is ejected. One dose is generally sufficient.

Another remedy, and a very simple yet effectual one, is tobacco smoking. Place the chicks in a small box or lined hamper. The box should have a lid and have two or three ventilation holes at the top. Then having filled your pipe with strong tobacco, puff the clouds of smoke into the box or basket until you cannot see the chicks. This causes them to cough violently and expel the thread-like worms from the throat which are the cause

of all the mischief. Gapes is most prevalent where the soil is damp, and where chicks have access to water that has been exposed to the sun's rays, and the same applies to diphtheria which often breaks out after a spell of very wet weather.

On dry sandy soil and where the water is kept pure and sweet, Gapes is unknown.

TICKS

If your chicks mope about with wings drooping and a distressed appearance look for "Ticks." These are only to be found on the head and round the throat. They must be looked for with great care, as they bury their heads in the skin of the chick and suck the blood, the loss of the vital fluid causing the chick to droop and die.

Treatment: apply to the roots of the feathers on head, neck, and throat, Carbolic Sweet Oil.

Another remedy is to mix one part paraffin to three parts *boiling* water. The paraffin should be added when the water is boiling. This is important. The head and round the throat should be just touched with this mixture after it has been allowed to cool, but not before, and it must *not* be rubbed in the feathers but simply touch the head and upper part of throat with it, using a piece of linen rag for the purpose.

Lice are a nuisance to poultry, and from which our Bantams unfortunately have no immunity. A good dust bath is a grand preventative, and especially if a little sulphur or insect powder can be dusted over it. However, a good practice is to periodically go over the birds with Keating's insect powder, a safe and sure remedy against these parasites. Early summer is the worst time for them, and at such times one can hardly go wrong with the dusting box.

Feather Disease, or what is sometimes called feather-curl, very frequently troubles Bantams. It generally shows itself worst on the breast, where the feathers appear dry and shrivelled. Their abnormal appearance is caused by a parasite, which practically sucks the virtue out of them and thus causes them to curl. A good way of dealing with this trouble is to prepare a solution of paraffin with boiling water in the proportion of half and half. Rub this well into the feathers with a piece of rag; one dressing will usually be found to be sufficient.

Egg-bound.—This is the most serious trouble to which Bantams are liable. The sooner it is noticed and taken in hand the easier it is to cure. If the bird is noticed on the nest an unusual length of time it may at once be assumed that she is finding difficulty in laying. Do not wait until matters have got beyond remedy on the supposition that she will manage it by and by. It is safest to take her at once, before her strength is exhausted and rupture has occurred, and place her up to the thighs in water, as hot as you can bear your hand in, for ten minutes or a quarter of an hour, and then after drying her with a towel, grease the vent well with vaseline. If the egg is in view, a little gentle pressure from the outside with the thumb and fingers will often eject the egg; but it must be done very carefully, or a fracture of the egg will result, and then the case usually becomes fatal.

I have heard of a recent case where the sufferer was kept in the hot water for three-quarters of an hour before a change for the better ensued, the egg being in view the whole time, and the vent syringed with warm olive oil. But perseverance was rewarded in due time, and the egg produced a chick. The whole operation is in reality very simple, but requires great care and some patience.

Scaly-leg is another complaint from which Bantams sometimes suffer, and which is generally caused by very scaly-legged hens being used as brooders. The scale referred to is a parasite feeding on the leg, so the sooner it is taken in hand the better. Thoroughly soak the bird's legs in hot water then brush them well with soap and a hard nail brush; afterwards anoint them with a dressing of sulphur ointment, made of vaseline and flowers of sulphur.

In slight cases a thoroughly good wash, and a good rubbing with crude paraffin oil will suffice to get rid of the parasite.

These seem to be the principal diseases attacking Bantams, and if the young fancier faithfully carries out the instructions I have given for their treatment, no difficulty will be experienced in effecting a cure.

In conclusion, I may say that throughout this work my sole aim has been to assist the novice as far as I possibly can, and from the multitude of grateful letters that I received while the book in article form was running through *The Feathered World*, I derived the greatest pleasure to learn that my endeavours have been so heartily appreciated.

THE FEATHERED WORLD,

9, Arundel Street,

Strand, London, W.C.2.

SOME USEFUL BOOKS.

ANATOMY OF THE FOWL. By P. PROUD. With full-page Plates, showing the Organs of the Fowl, with details as to their functions. Post free 8d. (16 cents).

BANTAMS AS A HOBBY. By P. PROUD, revised by H. INMAN. Contains full descriptions of Game and Variety Bantams, and gives practical information as to Housing, Breeding and Exhibiting. Over Thirty Illustrations. Post free, paper 1s. 9d. (42 cents).

BRAHMAS AND COCHINS. Illustrated. Second Edition. By L. C. R. NORRIS-ELYE. The Standard work on these grand old breeds. Post free 1s. 3d. (30 cents).

BRITISH POULTRY AND POULTRY KEEPING. By S. H. LEWER. The superb handbook on breeds of poultry. Was specially prepared for the World's Congress, with 14 coloured and 185 other Illustrations. Post free, paper 1s. 3d. (30 cents), cloth 3s. (75 cents).

CAMPINE FOWL. By the Rev. and Mrs. HARVEY. An interesting work on this popular breed. Coloured Frontispiece and other Illustrations. Post free 8d. (16 cents).

CHICKENS, MANAGEMENT OF. By P. PROUD. Directions on Feeding from the Shell to Maturity, also notes on the sitting hen and chicken ailments. Post free 4d. (8 cents).

DISEASES OF POULTRY. By PROF. WOODROFFE HILL. Revised and enlarged, with a Dictionary of Diseases and their Symptoms. Invaluable to every Poultry Fancier. Post free 1s. 3d. (30 cents).

GAME FOWL. By P. PROUD. Besides giving full information on the present-day Old English and Modern Game, includes a reprint of "The Cocker," written in 1815, which contains much advice applicable to-day. Illustrated. Post free, paper 1s. 10d. (50 cents), cloth 3s. 4d. (80 cents).

GEESE AND TURKEYS. By P. PROUD. Deals with all the principal varieties of both breeds, with hints on Hatching, Feeding, Fattening, etc. Post free 7d. (14 cents).

HAMBURGH UP TO DATE. By CHARLES HOLT. Illustrated, and contains full information on How to Breed all the Varieties; also Standards. Post free 1s. 3d. (30 cents).

INCUBATION, NATURAL AND ARTIFICIAL. By J. H. SUTCLIFFE. The Standard work on Hatching and Rearing. Fully Illustrated. Post free 1s. 9d. (42 cents).

INCUBATORS, MANAGEMENT OF. By P. PROUD. Also gives details on how to use Rearingers successfully. Post free 4d. (8 cents).

INTENSIVE POULTRY CULTURE. By Mrs. BAYNES. The practical experience of one who has proved its value. Fully illustrated. Post free 1s. (24 cents).

LEGHORNS. By L. C. VERREY. Two Coloured Plates, many Illustrations. Deals with all varieties of this popular breed, with full details as to mating, treatment, etc. Post free, 1s. 9d. (42 cents) paper, 2s. 9d. (66 cents) cloth.

MINORCA, THE. By T. H. HARRISON. Illustrated. An interesting treatise by an old fancier of the Variety. Post free 1s. 3d. (30 cents).

MODERN MINORCA FOWL, THE. By FRED TOOTILL. With Notes of Special Interest on Management and Preparing for Exhibition. Post free 1s. 3d. (30 cents).

OLD ENGLISH GAME FOWL. By HERBERT ATKINSON. Illustrated. The Standard work on this grand old breed. Post free 1s. 9d. (42 cents).

ORPINGTON, THE. By E. CAMPBELL. Deals with Mating for Colour, etc. Feeding and Management. With new chapters on White, Jubilee, Spangled, Blue and Cuckoo, by WILLIAM H. COOK. Gives Standards. Illustrated. Post free 1s. 3d. (30 cents).

PLYMOUTH ROCK FOWL. By Dr. J. P. CARTWRIGHT. The latest and fullest descriptive work on all varieties, with two Coloured Plates and several Illustrations. Post free 1s. 9d. (42 cents).

POULTRY KEEPERS' TROUBLES. By H. INMAN. In this book Mr. Inman has compiled from the queries sent him a set of helpful hints which we hope may solve some of the difficulties met with by Poultry-keepers. Post free 1s. 2d. (28 cents).

POULTRY YARD RECORD. An Account Book in which all details connected with Fowls may be readily entered. Post free 1s. 4d. (32 cents).

POULTRY MANAGEMENT THROUGHOUT THE YEAR. By CLEM WATSON. An Indispensable Guide, month by month, to the Poultry-keeper. Post free 1s. 3d. (30 cents).

HANDBOOKS ON POULTRY. By E. COBB.
Breeding for Exhibition. Post free 8d. (16 cents).
Feeding and Rearing of Chickens. Post free 8d. (16 cents).
Housing and Management of Stock. Post free 8d. (16 cents).
Incubation, Artificial and Natural. Post free 8d. (16 cents).
Preparing Poultry for Exhibition. Post free 8d. (16 cents).
Sick Fowls and their Treatment. Post free 8d. (16 cents).
Fattening Fowls. Post free 1s. 2d. (26 cents).

PRACTICAL POULTRY KEEPING FOR SMALL-HOLDERS. By J. STEPHEN HICKS and WILFRED H. G. EWART. A well-illustrated Handbook, full of practical information for Poultry-keepers generally, and with special reference to Poultry Management on Small-holdings. Post free 8d. (16 cents).

RHODE ISLAND RED, THE. By Miss M. H. CLAY. A new and deeply interesting work on the breed by one of its most skilled breeders. Illustrated. Post free 1s. 3d. (30 cents).

SCIENTIFIC BREEDING AND FEEDING OF POULTRY. By the late ALEX COMYNS, B.A., LL.B. Post free 10d. (20 cents).

SEBRIGHT BANTAM, THE. By Dr. WISEMAN CUNNINGHAM. Illustrated. Post free 1s. 2d. (28 cents).

STOCK AND EXHIBITION POULTRY BREEDING RECORD. By S. DE M. CAREY. Specially prepared for Fancy and Utility Poultry-keepers. Post free 11d. (22 cents).

SUBURBAN POULTRY-KEEPING. By T. H. HARRISON. Post free, paper 1s. 3d. (30 cents), cloth 2s. 4d. (56 cents).

TABLE-CHICKEN REARING. Intensive and Semi-Intensive. By F. HESKETH. Illustrated. 8d., post free (16 cents).

WATERFOWL, FANCY. By FRANK FINN, F.Z.S. Treats of Ducks, Geese and Swans of the most easily-obtained Varieties. Profusely Illustrated. Post free 3s. 4d. (80 cents), in cloth only.

WYANDOTTE, THE. By H. P. RAINES and J. P. W. MARX. Seventh Edition, revised by J. H. BROOKSBANK. Treats of all varieties, with the Club Standards. Coloured Plates and Numerous Illustrations. Post free 1s. 9d. (42 cents).

Also Books on Pigeons and Cagebirds, and a wide assortment of Coloured Plates and postcards of poultry, pigeons, canaries and British birds, list on application to

THE FEATHERED WORLD,

9, ARUNDEL STREET, STRAND, LONDON, W.C. 2.

